

Checklist of natural enemies of *Lipaphis erysimi* (Kaltenbach, 1843) and *Lipaphis pseudobrassicae* (Davis, 1914) (Hemiptera: Aphididae) and their host plant associations in India

Rajendra Singh 

Department of Zoology, Deendayal Upadhyay University of Gorakhpur, U.P., India

*Corresponding author ✉: rsinghpu@gmail.com

Citation: Singh, R. (2024). Checklist of natural enemies of *Lipaphis erysimi* (Kaltenbach, 1843) and *Lipaphis pseudobrassicae* (Davis, 1914) (Hemiptera: Aphididae) and their host plant associations in India. *Journal of Animal Diversity*, 6 (3): 1–25. <http://dx.doi.org/10.22034/JAD.2024.6.3.1>

Abstract

The present study summarizes the predators and parasitoids of two species, the mustard aphid *Lipaphis erysimi* (Kaltenbach, 1843) and the turnip aphid *Lipaphis pseudobrassicae* (Davis, 1914) infesting various food plants distributed in various states and union territories of India. These predators belong to four orders of the class Insecta: Coleoptera (Coccinellidae), Diptera (Chamaemyiidae, Syrphidae), Hemiptera (Anthocoridae, Geocoridae, Reduviidae) and Neuroptera (Chrysopidae, Hemerobiidae), and one order of class Arachnida, the order Araneae (spiders) (6 families). A total of 100 species of predators were observed to feed on the mustard aphid on 23 food plant species distributed in 26 states/union territory of India. The greatest numbers of predators of both species belong to the families Coccinellidae (47 species) and Syrphidae (27 species). Four species of predatory bugs (Hemiptera), seven species of lacewings (Neuroptera) and fourteen species of spiders are reported to prey on *Lipaphis erysimi* in India. Most of the tritrophic associations (triplets, predators–preys–host plants) of these predators are reported from Uttar Pradesh (98 triplets) followed by West Bengal (58 triplets), Punjab (53 triplets), Manipur (48 triplets), Tripura (40 triplets) and other states/union territories (less than 40 triplets). A total of 16 species of predators belonging to Coccinellidae, Syrphidae and Chrysopidae are reported to prey on *Lipaphis pseudobrassicae* only in seven states/union territory of India, mostly in Jammu and Kashmir. Two families of the parasitoids, Aphelinidae and Braconidae (subfamily Aphidiinae) parasitize *L. erysimi* in India. Aphelinidae is represented by 5 species distributed in seven states/union territory and Braconidae by 17 species on 20 host plants in 24 states/union territory of India. Most of the tritrophic associations (triplets, parasitoids–hosts–host plants) of these parasitoids are reported from Manipur (15 triplets) followed by Uttar Pradesh (11 triplets) and less than 10 triplets in other states/union territories. Only two identified species of Braconidae were observed to parasitize *L. pseudobrassicae* in India on two host plant species distributed only in Jammu and Kashmir, Karnataka and Punjab.

Editor-in-Chief: Dr. Ali Gholamifard

Assistant Editor: Dr. Erica DeMilio

Subject Editors: Professor Oldřich Nedvěd
Dr. John Caleb

Received: 8 August 2024

Revised: 29 August 2024

Accepted: 22 September 2024

Published online: 30 September 2024

Key words: Aphelinidae, biocontrol, Braconidae, Chrysopidae, Coccinellidae, mustard aphid, parasitoids, predators, spiders, Syrphidae

Introduction

Lipaphis (*Lipaphis*) *erysimi* (Kaltenbach, 1843) (Hemiptera: Aphididae), commonly known as mustard

aphid, is a polyphagous sucking insect pest and is associated with 52 species of host plants belonging to 42 genera in 19 families in India (Singh and Singh, 2017; Singh et al., 2023; Tiwari et al., 2024a) of which

several plant species are highly economically important. It is considered a notorious and key pest of several brassica crops, particularly oil seed crops (rapeseed mustard, yellow mustard, black mustard) (Bakhtia and Sekhon, 1989) and vegetable crops (*Brassica oleracea* L. var. *botrytis*, *B. o.* var. *capitata*, *B. rapa* L., *Raphanus sativus* L.) (Shukla et al., 1994). Another allied species of the genus is *Lipaphis* (*Lipaphis*) *pseudobrassicae* (Davis, 1914), commonly known as turnip aphid, which is a little-known species feeding mostly on brassica plants along with the mustard aphid. In mustard fields, sometimes nothing remains to harvest after their infestation (Palial et al., 2022). Both species infest mostly the inflorescences, buds, and young leaves and stems of its host plants. Both nymphs and adults (wingless and winged forms) suck the sap of the host plant, which causes wilting and flower dropping. Infested parts are deformed and crinkled, which hinders plant growth. In addition to direct damage, the aphids also excrete a large amount of honeydew, on which a black fungus called sooty mould develops and interferes with the normal functioning of the photosynthetic activity of the plant (Singh, 2014a). In addition, like other aphid species, *L. erysimi* also transmits several viral diseases, e.g., cabbage black ringspot virus, radish mosaic virus, cauliflower mosaic virus and turnip mosaic virus (Capinera, 2001). Because of its pest status, several practices have been applied to manage the population of *L. erysimi* in fields such as cultural, biological and chemical techniques (Sachan, 1997; Mishra and Gaurav, 2022).

It is an established fact that anthropogenic activities have significantly reduced global biodiversity (Prodanova et al., 2023). Therefore, one of the most significant challenges facing our earth today is its conservation, which includes the information about availability of fauna and flora in a given ecosystem. Insects are one of important elements of the species diversity. They are not only pests but also play a critical role in pollination, maintaining soil structure and fertility, seed dispersal, nutrient cycling, producing useful compounds, scavenging, providing food for other animals and managing pest insect populations by parasitizing and consuming them (Singh, 2014a). The natural enemies (parasitoids and predators) of insect pests have an essential role in maintaining their population in equilibrium and thus their balance in the ecosystem. The importance of conserving the natural enemies of insect pests is globally emphasized as these insects not only act as bioagents in controlling the pest population naturally, but are also used in classical and conservation biocontrol programs (Heimpel and Mills, 2017).

Among the aphidophagous arthropods, ladybird beetles (Coleoptera: Coccinellidae) stand first due to their species diversity and having high foraging and predatory efficiencies (Pervez et al., 2020a). Other aphidophagous arthropods are the larvae of syrphid flies (Diptera: Syrphidae) (Joshi and Ballal, 2013; Joshi et al., 2023;

Ahmad and Kumari, 2024), lacewings (Neuroptera: Chrysopidae, Hemerobiidae) (Bakthavatsalam and Varshney, 2023; Singh et al., 2024a), spiders (Arachnida: Araneae) (Singh et al., 2024b), and some predatory bugs (Ballal et al., 2023). Among the parasitoids, only members of two families, Braconidae and Aphelinidae (Hymenoptera) are reported to parasitize aphids (Singh and Singh, 2016).

The main objective of this compilation is to prepare a checklist of natural enemies of *Lipaphis erysimi* and *Lipaphis pseudobrassicae* and their food plant associations across the country that may help in their natural conservation, aiming to reduce the pest population and to protect the crops infested by them. Some of them, e.g., *Coccinella septempunctata* Linnaeus (a predator, Shenhmar and Brar, 1995) and *Diaeretiella rapae* (M'Intosh, 1855) (a parasitoid, Rajput, 2022) have been successfully utilized in its biological control on brassica crops in India. Most of the natural enemies recorded to prey or parasitize *Lipaphis erysimi* are polyphagous.

The present checklist is based on the primary data of published literature on the natural enemies of *L. erysimi* and *L. pseudobrassicae* extracted from books, book chapters, journals, proceedings of conferences, Records and Fauna Series of the Zoological Survey of India, Kolkata, and a few authentic theses available on Shodhganga (<https://shodhganga.inflibnet.ac.in>) up to August 15, 2024. Records of aphidophagous arthropods that do not name their specific prey species were excluded from this checklist. In most of the recent previous literature, there are several errors in the scientific names of the insects, both predators as well as their preys (aphids), and food plants because of their modified status and other nomenclatural decisions and clarifications. The names of aphids, as well as plants that were misspelt in the original records have been corrected where we logically ascertained the intended species. In the present checklist, attempts have been made to provide the valid scientific names of the predators following GBIF (2024), WSC (2024), with aphids following Favret (2024) and for plants, following (WFO, 2024). For detailed synonymy of the valid species, the above references should be consulted.

The district map of the Indian states used in Figures 1–3 was taken from the website <https://www.mapsofindia.com> and numbers of triplets (predator–aphid prey–food plant) recorded in each state/union territories are illustrated.

Results

Predatory arthropods of *Lipaphis erysimi* mostly belong to four orders of the class Insecta: Coleoptera, Diptera, Hemiptera and Neuroptera and one order of class Arachnida: Araneae (spiders) (6 families). A total 100 species of predators were observed to prey the mustard aphid on 23 food plant species distributed in 26 states/union territory of India (Table 1).

Table 1: Number of *Lipaphis erysimi* predator species with numbers of host plant species, associations and presence in Indian state/union territories for each higher taxon.

Class	Order	Families of predators	Number of predator species	Number of host plant species	Number of predator-prey-host plant associations	Number of states/union territories
Arachnida	Araneae	1. Araneidae	2	1	2	2
		2. Lycosidae	2	3	6	1
		3. Oxyopidae	1	2	2	1
		4. Salticidae	4	5	9	2
		5. Theridiidae	1	1	1	1
		6. Thomisidae	4	4	7	2
		Subtotal	14	5	27	3
Insecta	Coleoptera	1. Coccinellidae	47	19	154	25
	Diptera	1. Chamaemyiidae	1	1	1	3
		2. Syrphidae	27	15	87	22
		Subtotal	28	15	88	22
	Hemiptera	1. Anthocoridae	2	2	4	1
		2. Geocoridae	1	3	3	1
		3. Reduviidae	1	1	1	1
		Subtotal	4	5	8	3
	Neuroptera	1. Chrysopidae	4	8	15	13
		2. Hemerobiidae	3	5	7	5
		Subtotal	7	8	22	14
Total			100	23	299	26

**Figure 1:** Map showing the number of tritrophic associations (predator-aphid host-food plant) of aphidophagous arthropods preying on *Lipaphis erysimi* in different states and union territories of India. <https://www.mapsofindia.com>




Figure 2: Map showing the number of tritrophic associations of aphidophagous ladybird beetles preying on *Lipaphis erysimi* in different states and union territories of India. <https://www.mapsofindia.com>




Figure 3: Map showing the number of tritrophic associations of aphidophagous hover flies preying on *Lipaphis erysimi* in different states and union territories of India. <https://www.mapsofindia.com>

The observed predator species comprise 14 species of spiders, 47 species of ladybird beetles (Coleoptera: Coccinellidae), 1 species of Chamaemyiidae and 27 species of Syrphidae (Diptera), 2 species of Anthocoridae, 1 species of Geocoridae and 1 species of Reduviidae (Hemiptera), and 4 species of Chrysopidae and 3 species of Hemerobiidae (Neuroptera). Most of the tritrophic associations (triplets, predators–preys–host plants) of these predators were reported from Uttar Pradesh (98 triplets) followed by West Bengal (58 triplets), Punjab (53 triplets), Manipur (48 triplets), Tripura (40 triplets) and other states/union territories (less than 40 triplets) (Fig. 1). The actual number of species can be slightly lower due to identification of some species reported in literature only to genus level, but here treated as separate species.

Checklist of predators of *Lipaphis erysimi* infesting food plants in India

A. Class: Arachnida; Order: Araneae

The order Araneae includes spiders, which are an important element of the earth's ecosystem. Being predatory, they consume mostly insects, keeping their population under control and hardly pose any danger to humans (Nyffeler and Birkhofer, 2017). However, unlike insect predators, their potential as biocontrol agents is less exploited to its fullest. Recently, Singh et al. (2024b) reported a total of 77 species of aphidophagous spiders in India preying on 53 species of aphids infesting 58 species of plants. Herewith, 14 species of spiders are listed belonging to six families that prey on the mustard aphid in only three states of India. Indeed, very limited attempts have been made to record these aphidophagous spiders in India and an intensive and extensive survey plan to record them in unexplored areas is required.

Family 1: Araneidae

Eriovixia sp.

Brassica rapa L. – Maharashtra (Kale et al., 2020)

Neoscona sp.

Brassica rapa L. – Uttarakhand (Debnath, 1991); Maharashtra (Kale et al., 2020)

Family 2: Lycosidae

Pardosa pseudoannulata (Bösenberg and Strand, 1906)

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Wadicosa fidelis (O. Pickard-Cambridge, 1872)

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Family 3: Oxyopidae

Oxyopes javanus Thorell, 1887

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica sp. – West Bengal (Agarwala et al., 1981a)

Family 4: Salticidae

Hasarius adansoni (Audouin, 1826)

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Marpissa sp.

Brassica rapa L. – Uttarakhand (Debnath, 1991)

Phintella vittata (C. L. Koch, 1846)

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica sp. – West Bengal (Agarwala et al., 1981a)

Plexippus paykulli (Audouin, 1826)

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Family 5: Theridiidae

Theridion sp.

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Family 6: Thomisidae

Camaricus formosus Thorell, 1887

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica rapa L. – West Bengal (Agarwala et al., 1981a)

Thomisus projectus Tikader, 1960

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Thomisus pugilis Stoliczka, 1869

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Thomisus sp.

Brassica rapa L. – Maharashtra (Kale et al., 2020)

Class: Insecta; Order: Coleoptera

Family: Coccinellidae

These are commonly known as ladybird beetles. Though several species of this family are major agricultural pests, at least 260 species are insectivorous, feeding on soft insects like aphids, scale insects, mealy bugs, whiteflies, etc., in India (Omkar and Pervez, 2004). In this checklist, 47 species of coccinellids visiting 19 species of plants are reported, feeding on *Lipaphis erysimi* in 25 states/union territories of India (Table 1). Most of the tritrophic associations (triplets, predators–preys–host plant) of these ladybird beetles are reported from Uttar Pradesh (72 triplets) followed by Manipur (26 triplets), Punjab (25 triplet), West Bengal (23 triplets), Jammu and Kashmir (22 triplets), Uttarakhand (20 triplets), and less than 20 triplets in other states/union territories (Fig. 2). More than 60% species of these ladybird beetles prey on the mustard

aphid on oilseed brassica crops (*Brassica juncea* L., *Brassica napus* L., *Brassica nigra* (W.D.J. Koch, *Brassica rapa* L.) followed by brassica vegetables (*Brassica oleracea* L. – cabbage, cauliflower), and radish (*Raphanus sativus* L.). A checklist of ladybird beetles feeding on *Lipaphis erysimi* infesting food plants in different states of India follows:

***Adalia bipunctata* (Linnaeus, 1758)**

Brassica napus L. – Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Chaudhary and Singh, 2012; Tiwari et al., 2024b)

***Adalia tetraspilota* (Hope, 1831)**

Brassica napus L. – Manipur (Chakrabarti et al., 2012)

Brassica rapa L. – Jammu and Kashmir (Bali, 1987; Mir, 2013)

***Anegleis cardoni* (Weise, 1892)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Singh, 2009)

Brassica rapa L. – Uttar Pradesh (Afroze, 1999; Omkar et al., 2009)

Brassica sp. – Manipur (Chakrabarti et al., 2012); Uttarakhand (Pal and Bhatt, 2018)

***Brumoides suturalis* (Fabricius, 1798)**

Brassica juncea (L.) Czern. – Assam (Das, 2020); Punjab (Soni et al., 2004); Uttar Pradesh (Tiwari et al., 2024b); Uttarakhand (Sharma and Joshi, 2010); West Bengal (Ghosh, 1983)

Brassica rapa L. – Himachal Pradesh (Gautam, 1990); Uttar Pradesh (Tiwari et al., 2024b)

***Brumus* sp.**

Brassica carinata A. Braun – Punjab (Kumar, 2015)

Brassica juncea (L.) Czern. – Punjab (Kumar, 2015)

Brassica napus L. – Punjab (Kumar, 2015)

Brassica nigra W.D.J. Koch – Punjab (Kumar, 2015)

Brassica rapa L. – Punjab (Kumar, 2015)

Eruca vesicaria (L.) Cav. – Punjab (Kumar, 2015)

***Calvia punctata* (Mulsant, 1853)**

Brassica rapa L. – Jammu and Kashmir (Khan et al., 2009; Mir, 2013)

***Cheilomenes sexmaculata* (Fabricius, 1781)**

Arivela viscosa (L.) Raf. – Kerala (Thamilvel, 2009)

Brassica carinata A. Braun – Punjab (Kumar, 2015)

Brassica juncea (L.) Czern. – Assam (Das, 2020); Chhattisgarh (Ray, 1967; Kolhekar et al., 2019); Delhi (Pandi et al., 2012); Gujarat (Patel, 2015); Haryana (Kalra, 1988); Himachal Pradesh (Soni et al., 2021); Jammu and Kashmir (Mir, 2013); Karnataka (Joshi et al., 1999a); Madhya Pradesh (Mishra and Kanwat, 2018); Manipur (Rajeshwari and Singh, 2022); Meghalaya (Damitre, 2019); Punjab (Sharma et al., 1997; Soni et al., 2004); Rajasthan (Singh et al., 2009; Ram et al., 2022); Tripura (Datta, 2020); Uttar Pradesh (Shukla and Kumar,

2024; Tiwari et al., 2024b); Uttarakhand (Pal and Bhatt, 2018; Verma et al., 2023); West Bengal (Ghosh, 1983)

Brassica napus L. – Manipur (Shantibala, 1989); Punjab (Kumar, 2015); Uttar Pradesh (Tiwari et al., 2024b)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998)

Brassica oleracea L. var. *capitata* – Andhra Pradesh (Rao et al., 1997); Gujarat (Varmora et al., 2009; Chakraborty, 2012); Odisha (Mandal and Patnaik, 2006); Manipur (Bijaya et al., 1996); Maharashtra (Shaikh et al., 2020); Nagaland (Waluniba et al., 2016); Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Bihar (Prabhakar and Roy, 2010; Kumar et al., 2016); Gujarat (Chakraborty, 2012); Haryana (Rana, 2006); Jammu and Kashmir (Mir, 2013; Bhat, 2017); Madhya Pradesh (Verma et al., 1983); Maharashtra (Kale et al., 2020; Shaikh et al., 2020); Odisha (Parija and Biswal, 2023); Punjab (Kumar, 2015; Shenhmar and Brar, 1995); Rajasthan (Sharma, 1973; Singh et al., 2008); Uttar Pradesh (Chaudhary and Singh, 2012; Tiwari et al., 2024b); Uttarakhand (Pervez and Chandra, 2018)

Brassica sp. – Jammu and Kashmir (Khan et al., 2009)

Eruca vesicaria (L.) Cav. – Punjab (Kumar, 2015)

Raphanus sativus L. – Kerala (Thamilvel, 2009); Maharashtra (Shaikh et al., 2020); Uttarakhand (Pervez and Chandra, 2018)

Vigna unguiculata (L.) Walp. – Andhra Pradesh (Rao et al., 1997)

Unknown plant sp. – Tripura (Agarwala et al., 1987)

***Chilocorus infernalis* Mulsant, 1853 [syn. *Chilocorus bijugus* Mulsant, 1853]**

Raphanus sativus L. – Himachal Pradesh (Sharma et al., 2015)

***Chilocorus nigritus* (Fabricius, 1798)**

Brassica juncea (L.) Czern. – Uttarakhand (Sharma and Joshi, 2010)

Brassica rapa L. – Uttar Pradesh (Omkar and Bind, 1995)

***Coccinella californica* Mannerheim, 1843**

Brassica rapa L. – Odisha (Parija and Biswal, 2023)

***Coccinella luteopicta* (Mulsant, 1866)**

Raphanus sativus L. – Himachal Pradesh (Sharma et al., 2015)

***Coccinella magnopunctata* Rybakow, 1889**

Brassica juncea (L.) Czern. – Jammu and Kashmir (Mir, 2013)

***Coccinella repanda* Thunberg, 1781**

Brassica juncea (L.) Czern. – Uttar Pradesh (Kishor et al., 2023; Tiwari et al., 2024b)

Brassica juncea (L.) Czern. – Assam (Saharia, 1981)

Brassica oleracea L. var. *capitata* – Chhattisgarh (Singh, 2014b)

Brassica rapa L. – Punjab (Srivastava et al., 1982); Uttar Pradesh (Tiwari et al., 2024b)

Coccinella septempunctata* Linnaeus, 1758Brassica carinata* A. Braun – Punjab (Kumar, 2015)*Brassica juncea* (L.) Czern. – Assam (Das, 2020); Bihar (Mishra and Mukharjee, 2020); Chhattisgarh (Kashyap et al., 2018; Patel et al., 2004); Delhi (Shankarganesh et al., 2015); Gujarat (Kulkarni and Patel, 2001; Patel et al., 2019); Haryana (Kalra, 1988; Singh and Singh, 1994); Himachal Pradesh (Soni et al., 2021); Karnataka (Joshi et al., 1999a); Madhya Pradesh (Lal et al., 2018; Mishra and Kanwat, 2018); Manipur (Chitra Devi et al., 1997, 2002); Meghalaya (Manpoong et al., 2016; Damitre, 2019); Punjab (Sharma et al., 1997; Kumar, 2015); Rajasthan (Chand et al., 2022; Yadav et al., 2023); Tripura (Gangopadhyay, 1980; Datta, 2020); Uttar Pradesh (Singh, 2013; Tiwari et al., 2024b); West Bengal (Ghosh, 1983)*Brassica napus* L. – Manipur (Shantibala, 1989); Punjab (Kumar, 2015); Uttar Pradesh (Tiwari et al., 2024b)*Brassica nigra* W.D.J. Koch – Jammu and Kashmir (Salalia and Isher, 2011); Manipur (Shantibala, 1989); Uttar Pradesh (Tiwari et al., 2024b); West Bengal (Ghosh et al., 1981)*Brassica oleracea* L. – Uttarakhand (Ghosh et al., 1991)*Brassica oleracea* L. var. *botrytis* – Bihar (Ahmad et al., 2012); Kerala (Ravi et al., 2014); Manipur (Devjani and Singh, 1998); Rajasthan (Choudhary et al., 2023); Uttar Pradesh (Tiwari et al., 2024b)*Brassica oleracea* L. var. *capitata* – Chhattisgarh (Singh, 2014b); Gujarat (Varmora et al., 2009; Chakraborty, 2012); Kerala (Ravi et al., 2014); Maharashtra (Shaikh et al., 2020); Manipur (Bijaya et al., 1996); Nagaland (Waluniba et al., 2016); Odisha (Mandal and Patnaik, 2006); Punjab (Atwal and Sethi, 1963); Rajasthan (Bana et al., 2012); Uttar Pradesh (Tiwari et al., 2024b); West Bengal (Gurung et al., 2019)*Brassica oleracea* L. var. *italica* – Mizoram (Boopathi et al., 2020)*Brassica rapa* L. – Bihar (Ahmad et al., 2012; Goswami et al., 2016); Delhi (Shankarganesh et al., 2015); Gujarat (Zalavadia and Kapadia, 2000); Haryana (Rana, 2006); Himachal Pradesh (Lakhanpal and Raj, 1998); Jammu and Kashmir (Khan et al., 2009; Mir, 2013); Maharashtra (Panchal et al., 2013; Shaikh et al., 2020); Manipur (Devjani and Singh, 2006); Meghalaya (Dwivedi et al., 2021); Punjab (Atwal et al., 1971; Shenhmar and Brar, 1995); Rajasthan (Gour and Pareek, 2003; Singh and Meghwal, 2010); Uttar Pradesh (Hugar et al., 2008; Chaudhary and Singh, 2012); Uttarakhand (Bisht et al., 2001; Pervez et al., 2020b); West Bengal (Gurung et al., 2019; Maji et al., 2023)*Brassica* sp. – Uttarakhand (Pal and Bhatt, 2018)*Eruca vesicaria* (L.) Cav. – Punjab (Kumar, 2015)*Ranunculus* sp. – Bihar (Ahmad et al., 2012)*Raphanus sativus* L. – Maharashtra (Shaikh et al., 2020); Manipur (Shantibala, 1989); Tripura (Gangopadhyay, 1980); Uttar Pradesh (Tiwari et al., 2024b); Uttarakhand (Ghosh et al., 1991)***Coccinella transversalis* Fabricius, 1781***Arivela viscosa* (L.) Raf. – Kerala (Thamilvel, 2009)*Brassica carinata* A. Braun – Punjab (Kumar, 2015)*Brassica juncea* (L.) Czern. – Assam (Das, 2020; Pradhan et al., 2020a); Gujarat (Jadhav and Shukla, 2015); Karnataka (Joshi et al., 1999a); Madhya Pradesh (Mishra and Kanwat, 2018); Manipur (Chitra Devi et al., 2002; Rajeshwari and Singh, 2022); Meghalaya (Damitre, 2019); Punjab (Kumar, 2015); Tripura (Gangopadhyay, 1980; Ghosh et al., 2017); Uttar Pradesh (Kishor et al., 2023; Tiwari et al., 2024b); West Bengal (Ghosh, 1983)*Brassica napus* L. – Manipur (Shantibala, 1989); Punjab (Kumar, 2015); Uttar Pradesh (Tiwari et al., 2024b)*Brassica nigra* W.D.J. Koch – Manipur (Shantibala, 1989); Uttar Pradesh (Tiwari et al., 2024b); West Bengal (Ghosh et al., 1981)*Brassica oleracea* L. var. *botrytis* – Bihar (Ahmad et al., 2012; Prakash and Rani, 2015); Manipur (Bijaya et al., 1996; Devjani and Singh, 1998); Odisha (Mandal and Patnaik, 2006); Uttar Pradesh (Tiwari et al., 2024b); West Bengal (Gurung et al., 2019)*Brassica oleracea* L. var. *capitata* – Gujarat (Varmora et al., 2009); Nagaland (Waluniba et al., 2016); Uttar Pradesh (Tiwari et al., 2024b)*Brassica oleracea* L. var. *italica* – Mizoram (Boopathi et al., 2020)*Brassica rapa* L. – Assam (Borah and Dutta, 2010a); Bihar (Prabhakar and Roy, 2010; Goswami et al., 2016); Manipur (Devjani and Singh, 2006); Odisha (Parija and Biswal, 2023); Punjab (Kumar, 2015); Tripura (Gangopadhyay, 1980); Uttar Pradesh (Hugar et al., 2008; Tiwari et al., 2024b); West Bengal (Gurung et al., 2019; Maji et al., 2023)*Eruca vesicaria* (L.) Cav. – Punjab (Kumar, 2015)*Raphanus sativus* L. – Kerala (Thamilvel, 2009); Tripura (Gangopadhyay, 1980); Uttar Pradesh (Tiwari et al., 2024b)***Coccinella undecimpunctata* Linnaeus, 1758***Brassica juncea* (L.) Czern. – Bihar (Rahman, 1940); Punjab (Rahman, 1940)*Brassica napus* L. – Jammu and Kashmir (Bhat, 2017)*Brassica oleracea* L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)*Brassica oleracea* L. var. *capitata* – Uttar Pradesh (Tiwari et al., 2024b)*Brassica rapa* L. – Jammu and Kashmir (Bhagat and Matta, 2002); Uttar Pradesh (Chaudhary and Singh, 2012; Tiwari et al., 2024b)*Lagenaria siceraria* (Molino) Standl. – Jammu and Kashmir (Bhat, 2017)***Coccinella* sp.***Brassica rapa* L. – Uttarakhand (Bisht et al., 2001)***Coelophora saucia* (Mulsant, 1850)***Brassica napus* L. – Manipur (Shantibala, 1989)*Brassica oleracea* L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Uttar Pradesh (Tiwarei et al., 2024b)

***Harmonia axyridis* (Pallas, 1773)**

Brassica juncea (L.) Czern. – Assam (Pradhan et al., 2020b)

***Harmonia dimidiata* (Fabricius, 1781)**

Brassica juncea (L.) Czern. – Himachal Pradesh (Soni et al., 2021); Meghalaya (Damitre, 2019)

***Harmonia eucharis* (Mulsant, 1853)**

Brassica juncea (L.) Czern. – Jammu and Kashmir (Mir, 2013)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica rapa L. – Jammu and Kashmir (Mir, 2013); Uttar Pradesh (Tiwarei et al., 2024b); Uttarakhand (Bisht et al., 2001; Pervez et al., 2020b)

***Harmonia octomaculata* (Fabricius, 1787)**

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Tiwarei et al., 2024b)

***Hippodamia variegata* (Goeze, 1777)**

Brassica juncea (L.) Czern. – Haryana (Kalra, 1988; Singh and Singh, 1994); Himachal Pradesh (Soni et al., 2021); Madhya Pradesh (Mishra and Kanwat, 2017); Uttar Pradesh (Tiwarei et al., 2024b)

Brassica napus L. – Jammu and Kashmir (Bhat, 2017); Uttar Pradesh (Tiwarei et al., 2024b)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica rapa L. – Jammu and Kashmir (Bhagat and Matta, 2002; Mir, 2013); Uttar Pradesh (Tiwarei et al., 2024b); Uttarakhand (Ghosh et al., 1991; Pervez et al., 2020c)

Brassica sp. – Uttarakhand (Pal and Bhatt, 2018)

Raphanus sativus L. – Uttar Pradesh (Tiwarei et al., 2024b)

***Hippodamia variegata doubledayi* (Mulsant, 1850)**

Brassica juncea (L.) Czern. – Jammu and Kashmir (Mir, 2013)

Brassica rapa L. – Uttar Pradesh (Chaudhary and Singh, 2012)

***Hippodamia* sp.**

Brassica rapa L. – Uttarakhand (Bisht et al., 2001)

***Illeis cincta* (Fabricius, 1798)**

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica oleracea L. var. *gongylodes* – Jammu and Kashmir (Mir, 2013)

Brassica rapa L. – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica sp. – Uttarakhand (Pal and Bhatt, 2018)

***Illeis indica* Timberlake, 1943**

Brassica juncea (L.) Czern. – Jammu and Kashmir (Mir, 2013)

***Machimus indianus* Ricardo, 1919**

Brassica rapa L. – Uttarakhand (Bisht et al., 2001)

***Megalocaria dilatata* (Fabricius, 1775)**

Brassica napus L. – Manipur (Shantibala, 1989)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Tiwarei et al., 2024b)

Raphanus sativus L. – Manipur (Shantibala, 1989)

***Micraspis allardi* (Mulsant, 1866)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 1999; Singh, 2009)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica sp. – Manipur (Chakrabarti et al., 2012); Uttarakhand (Pal and Bhatt, 2018)

***Micraspis discolor* (Fabricius, 1798)**

Brassica juncea (L.) Czern. – Assam (Das, 2020); Meghalaya (Damitre, 2019); Tripura (Datta, 2020); West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *italica* – Mizoram (Boopathi et al., 2020)

Brassica oleracea L. – Odisha (Mandal and Patnaik, 2006); West Bengal (Gurung et al., 2019)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwarei et al., 2024b)

Brassica oleracea L. var. *capitata* – Chhattisgarh (Singh, 2014b)

Brassica rapa L. – Bihar (Goswami et al., 2016); Tripura (Agarwala et al., 1988); Uttar Pradesh (Tiwarei et al., 2024b); West Bengal (Gurung et al., 2019; Maji et al., 2023); Bihar (Prabhakar and Roy, 2010)

***Micraspis vincta* (Gorham, 1895)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Singh, 2009; Omkar and Pervez, 2000)

Brassica oleracea L. – Andhra Pradesh (Rao et al., 1997)

Brassica rapa L. – Uttar Pradesh (Tiwarei et al., 2024b)

Vigna unguiculata (L.) Walp. – Andhra Pradesh (Rao et al., 1997)

***Nephus regularis* (Sicard, 1929)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 1999)

Brassica sp. – Manipur (Chakrabarti et al., 2012)

***Oenopia conglobata* (Linnaeus, 1758)**

Brassica rapa L. – Jammu and Kashmir (Bhat, 2017)

***Oenopia kirbyi* Mulsant, 1850**

Brassica juncea (L.) Czern. – Meghalaya (Damitre, 2019)

Brassica napus L. – Manipur (Shantibala, 1989)

Raphanus sativus L. – Uttar Pradesh (Tiwarei et al., 2024b)

***Oenopia sexareata* (Mulsant, 1853)**

Brassica juncea (L.) Czern. – Meghalaya (Damitre, 2019)

***Priscibrum uropygialis* (Mulsant, 1853)**

Brassica juncea (L.) Czern. – Jammu and Kashmir (Mir, 2013)

Brassica rapa L. – Jammu and Kashmir (Mir, 2013)
Raphanus sativus L. – Himachal Pradesh (Sharma et al., 2015)

***Propylea dissecta* (Mulsant, 1850)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Singh, 2009)
Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Mizoram (Boopathi et al., 2020)

Brassica rapa L. – Uttar Pradesh (Omkar and Mishra, 2005; Tiwari et al., 2024b); Uttarakhand (Pervez and Kumar, 2017)

Raphanus sativus L. – Uttar Pradesh (Pervez and Omkar, 2004)

***Propylea japonica* (Thunberg, 1781)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 1999; Tiwari et al., 2024b)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Chaudhary and Singh, 2012; Tiwari et al., 2024b)

Brassica sp. – Manipur (Chakrabarti et al., 2012)

***Propylea luteopustulata* (Mulsant, 1850)**

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – Uttar Pradesh (Tiwari et al., 2024b); West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Manipur (Shantibala, 1989)

Brassica rapa L. – Jammu and Kashmir (Khan et al., 2009); Uttar Pradesh (Tiwari et al., 2024b)

Raphanus sativus L. – Himachal Pradesh (Sharma et al., 2015)

Rosa sp. – West Bengal (Agarwala et al., 1980)

***Propylea quatuordecimpunctata* (Linnaeus, 1758)**

Zea mays L. – West Bengal (Chakrabarti et al., 2012)

***Pseudaspidimerus trinotata* (Thunberg, 1781)**

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Poddar and Ghosh, 1984)

***Psyllobora bisoetonotata* (Mulsant, 1850)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 1999)

***Novius ruficollis* (Mulsant, 1850)**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 1999)

***Scymnus (Pullus) pyrocheilus* Mulsant, 1853**

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica rapa L. – Bihar (Prabhakar and Roy, 2010)

***Scymnus fuscatus* Boheman, 1859**

Brassica juncea (L.) Czern. – Uttar Pradesh (Omkar and Pervez, 2000)

Brassica oleracea L. var. *botrytis* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Uttar Pradesh (Tiwari et al., 2024b)

***Scymnus* sp.**

Brassica juncea (L.) Czern. – Assam (Das, 2020); Uttar Pradesh (Omkar and Pervez, 1999); Uttarakhand (Ghosh et al., 1991)

II. Order: Diptera

Among Diptera, the hover flies (Syrphidae) are the most common aphid predators, being widely distributed in the Nearctic, Neotropical and Palaearctic regions. Most of the Syrphidae provide dual service in nature, as an adult they help in crop pollination (Joshi et al., 2023) and as larvae, they consume several soft-bodied insects such as aphids, scale insects, mealy bugs, whiteflies, thrips, etc., and thus form an integral part of natural control of these insects. Syrphidae is the largest family with 27 aphidophagous species feeding on *Lipaphis erysimi* infesting 15 species of host plants distributed in 22 states/union territory of India (Table 1). The highest number of species of these hover flies prey on aphids on oilseed mustard crops followed by vegetable brassica crops. Most of the tritrophic associations (triplet, predators–prey–host plant) of these hover flies are reported from Tripura (29 triplets) followed by Punjab (23 triplets), Pradesh (21 triplets), Manipur (20 triplets), and less than 20 triplets in other states/union territories (Fig. 3). The representation of another family, Chamaemyiidae, is very poor. The detailed food plant associations of these flies are given below:

Family 1: Chamaemyiidae

***Leucopis* sp.**

Brassica juncea (L.) Czern – Punjab (Sharma et al., 1997); West Bengal (Ghosh, 1983); Haryana (Kalra, 1988)

Family 2: Syrphidae

***Allograpta javana* (Wiedemann, 1824)**

Brassica juncea (L.) Czern – Tripura (Agarwala et al., 1989; Datta, 2020)

Brassica oleracea L. var. *botrytis* – Tripura (Agarwala et al., 1989)

Brassica oleracea L. var. *capitata* – Tripura (Agarwala et al., 1989)

Brassica rapa L. – Uttar Pradesh (Tiwari et al., 2024b)

Raphanus sativus L. – Tripura (Agarwala et al., 1989)

***Baccha* sp.**

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

***Betasyrphus isaaci* (Bhatia, 1933)**

Brassica juncea (L.) Czern. – Meghalaya (Manpoong et al., 2016); Bihar (Rahman, 1940)

Brassica rapa L. – Meghalaya (Dwivedi et al., 2021); Uttar Pradesh (Tiwari et al., 2024b); Uttarakhand (Bisht et al., 2006)

Unknown plant sp. – Tripura (Agarwala et al., 1987)

***Betasyrphus serarius* (Wiedemann, 1830)**

Brassica juncea (L.) Czern. – Manipur (Chitra Devi et al., 2002); Punjab (Sharma et al., 1997); Tripura (Agarwala et al., 1989; Datta, 2020)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998; Singh et al., 2002); Tripura (Agarwala et al., 1989)

Brassica oleracea L. var. *capitata* – Manipur (Agarwala et al., 1984; Bijaya et al., 1996); Meghalaya (Agarwala et al., 1984); Tripura (Agarwala et al., 1989); West Bengal (Agarwala et al., 1982)

Brassica rapa L. – Jammu and Kashmir (Bhat and Bhagat, 2017); Uttar Pradesh (Tiwari et al., 2024b)

Raphanus sativus L. – Tripura (Agarwala et al., 1989)

***Betasyrphus* sp.**

Brassica juncea (L.) Czern. – Karnataka (Chinnu et al., 2023)

***Dideopsis aegrota* (Fabricius, 1805)**

Unknown aphid sp. – Tripura (Agarwala et al., 1987)

***Episyrphus balteatus* (De Geer, 1776)**

Brassica carinata A. Braun – Punjab (Kumar, 2015)

Brassica juncea (L.) Czern. – Haryana (Kalra, 1988; Singh and Singh, 1994); Himachal Pradesh (Soni et al., 2021); Manipur (Chitra Devi et al., 1997; Devi et al., 2011); Punjab (Sharma et al., 1997; Kumar et al., 1988); Tripura (Gangopadhyay, 1980; Agarwala et al., 1989); Uttar Pradesh (Singh, 2013; Tiwari et al., 2024b); West Bengal (Ghosh, 1983)

Brassica napus L. – Punjab (Kumar, 2015)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998; Singh et al., 2002); Tripura (Agarwala et al., 1989); Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996); Tripura (Agarwala et al., 1989)

Brassica rapa L. – Assam (Borah and Dutta, 2010b); Jammu and Kashmir (Bhagat and Matta, 2002; Bhat, 2008); Manipur (Devjani and Singh, 2006); Odisha (Parija and Biswal, 2023); Punjab (Kumar, 2015); Tripura (Gangopadhyay, 1980); Uttar Pradesh (Singh and Singh, 2013; Tiwari et al., 2024b); Uttarakhand (Bisht et al., 2006)

Brassica sp. – West Bengal (Roy and Basu, 1977)

Eruca vesicaria (L.) Cav. – Punjab (Kumar, 2015)

Raphanus sativus L. – Tripura (Gangopadhyay, 1980; Agarwala et al., 1989)

***Episyrphus viridaureus* (Wiedemann, 1824)**

Brassica juncea (L.) Czern. – Meghalaya (Manpoong et al., 2016); Punjab (Kumar et al., 1989; Sharma et al., 1997); Tripura (Agarwala et al., 1989; Datta, 2020)

Brassica oleracea L. var. *botrytis* – Tripura (Agarwala et al., 1989)

Brassica oleracea L. var. *capitata* – Tripura (Agarwala et al., 1989)

Brassica rapa L. – Meghalaya (Dwivedi et al., 2021); Punjab (Kumar et al., 1987)

Raphanus sativus L. – Tripura (Agarwala et al., 1989)

***Eristalis tenax* (Linnaeus, 1758)**

Brassica juncea (L.) Czern. – Chhattisgarh (Kashyap et al., 2018)

***Eumerus albifrons* Walker, 1852**

Brassica oleracea L. var. *capitata* – Odisha (Mandal and Patnaik, 2006)

***Eupeodes bucculatus* (Rondani, 1857)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988; Sharma et al., 1997)

Brassica rapa L. – Punjab (Kumar et al., 1987)

***Eupeodes confrater* (Wiedemann, 1830)**

Brassica juncea (L.) Czern. – Manipur (Chitra Devi et al., 1996, 2002); Punjab (Kumar et al., 1988; Sharma et al., 1997); Tripura (Gangopadhyay, 1980; Datta, 2020); Uttar Pradesh (Singh, 2013); West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998; Singh et al., 2002); Tripura (Agarwala et al., 1989); Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996); Tripura (Agarwala et al., 1989); Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Jammu and Kashmir (Bhat, 2008; Bhat and Bhagat, 2017); Manipur (Chitra Devi et al., 1996); Punjab (Kumar et al., 1987); Tripura (Gangopadhyay, 1980); Uttar Pradesh (Chaudhary and Singh, 2012; Tiwari et al., 2024b)

Brassica sp. – West Bengal (Roy and Basu, 1977)

Raphanus sativus L. – Tripura (Gangopadhyay, 1980; Agarwala et al., 1989)

***Eupeodes corollae* (Fabricius, 1794)**

Brassica napus L. – Jammu and Kashmir (Bhat and Bhagat, 2017)

Brassica rapa L. – Jammu and Kashmir (Bhat, 2008; Bhat and Bhagat, 2017); Uttarakhand (Bisht et al., 2006)

***Eupeodes* sp.**

Brassica napus L. – Himachal Pradesh (Lakhanpal and Raj, 1998)

Brassica oleracea L. var. *capitata* – Nagaland (Waluniba et al., 2016)

***Ischiodon scutellaris* (Fabricius, 1805)**

Brassica juncea (L.) Czern. – Assam (Pradhan et al., 2020b); Delhi (Lal and Gupta, 1953); Gujarat (Chouhan et al., 2011; Patel et al., 2019); Himachal Pradesh (Soni et al., 2021); Karnataka (Joshi et al., 1999b); Madhya Pradesh (Mishra and Kanwat, 2017); Manipur (Chitra Devi et al., 2002; Rajeshwari and Singh, 2022); Punjab (Kumar et al., 1988; Sharma et al., 1997); Rajasthan (Chand et al., 2022; Yadav et al., 2023); Tripura (Datta, 2020; Agarwala et al., 1989); Uttar Pradesh (Ali et al., 2009; Singh, 2013); West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998; Singh et al., 2002); Tripura (Agarwala et al., 1989); Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996); Odisha (Mandal and Patnaik, 2006); Tripura (Agarwala et al., 1989)

Brassica oleracea L. var. *gongylodes* – Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Jammu and Kashmir (Bhat and Bhagat, 2017); Maharashtra (Kale et al., 2020); Manipur (Chitra Devi et al., 1996; Devjani and Singh, 2006); Punjab (Kumar et al., 1987); Uttar Pradesh (Singh and Singh, 2013; Tiwari et al., 2024b); Uttarakhand (Karnatak and Thorat, 2006)

Brassica sp. – West Bengal (Roy and Basu, 1977)

Raphanus sativus L. – Tripura (Agarwala et al., 1989)

***Melanostoma orientale* (Wiedemann, 1824)**

Brassica juncea (L.) Czern – Punjab (Sharma et al., 1997)

Brassica rapa L. – Uttarakhand (Bisht et al., 2006)

Sinapis spp. – Himachal Pradesh (Agarwala et al., 1984); Punjab (Agarwala et al., 1984); Sikkim (Agarwala et al., 1984)

***Melanostoma univittatum* (Wiedemann, 1824)**

Brassica rapa L. – Jammu and Kashmir (Ahmad and Bhat, 1986); Uttarakhand (Bisht et al., 2006)

Cucumis sativus L. – Uttarakhand (Bisht et al., 2006)

***Paragus crenulatus* Thomson, 1869**

Unknown plant sp. – Tripura (Agarwala et al., 1987)

***Paragus serratus* (Fabricius, 1805)**

Brassica juncea (L.) Czern. – Karnataka (Joshi et al., 1999b); Manipur (Chitra Devi et al., 2002); West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. var. *botrytis* – Manipur (Singh et al., 2002); Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Jammu and Kashmir (Bhat and Bhagat, 2017); Manipur (Devjani and Singh, 2006); Uttar Pradesh (Tiwari et al., 2024b)

Raphanus sativus L. – Uttar Pradesh (Tiwari et al., 2024b)

***Scaeva albomaculata* (Macquart, 1842)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Scaeva latimaculata* (Brunetti, 1923)**

Brassica rapa L. – Punjab (Kumar et al., 1987)

Raphanus sativus L. – Punjab (Sharma et al., 1997)

***Scaeva selenitica* (Meigen, 1822)**

Brassica rapa L. – Uttarakhand (Bisht et al., 2006)

Sinapis sp. – Delhi (Agarwala et al., 1984)

***Sphaerophoria indiana* Bigot, 1884**

Brassica juncea (L.) Czern – Punjab (Kumar et al., 1988; Sharma et al., 1997)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015); Manipur (Devjani and Singh, 1998; Singh et al., 2002); Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996)

Brassica rapa L. – Punjab (Kumar et al., 1987); Uttar Pradesh (Tiwari et al., 2024b); Uttarakhand (Debnath, 1991)

Sinapis sp. – Delhi, Uttar Pradesh, Punjab (Agarwala et al., 1984)

***Sphaerophoria scripta* (Linnaeus, 1758)**

Brassica juncea (L.) Czern. – Assam (Das, 2020); West Bengal (Ghosh, 1983)

Brassica napus L. – Jammu and Kashmir (Bhat and Bhagat, 2017)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981; Agarwala et al., 1984)

Brassica rapa L. – Jammu and Kashmir (Ahmad and Bhat, 1986); Uttar Pradesh (Tiwari et al., 2024b); Uttarakhand (Bisht et al., 2001)

***Sphaerophoria* sp.**

Brassica oleracea L. var. *capitata* – Nagaland (Waluniba et al., 2016)

Unknown plant sp. – Karnataka (Agarwala et al., 1984)

***Syrphus* sp.**

Brassica juncea (L.) Czern. – Assam (Das, 2020)

Brassica oleracea L. var. *capitata* – Chhattisgarh (Singh, 2014b)

Brassica rapa L. – Jammu and Kashmir (Bhagat and Lone, 1984); Uttarakhand (Bisht et al., 2006)

***Xanthogramma* sp.**

Brassica oleracea L. – Rajasthan (Choudhary et al., 2023)

Brassica sp. – Delhi (Agarwala et al., 1984)

Coriandrum sp. – Delhi (Agarwala et al., 1984)

Order: Hemiptera

The order Hemiptera includes at least seven families of predatory bugs preying on aphids, scale insects, mealybugs, bugs, leaf hoppers, thrips, mites, caterpillars, etc., in India (Chellappan and Ranjith, 2023). However, only four hemipteran species belonging to three families (Anthocoridae, Geocoridae and Reduviidae) are reported as predators of *Lipaphis erysimi* in only West Bengal (Table 1) as mentioned below:

Family 1: Anthocoridae***Bilia castanea* (Carvalho, 1951)**

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981; Saxena, 1981)

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

***Orius albidipennis* (Reuter, 1884)**

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Family 2: Geocoridae***Geocoris ochropterus* (Fieber, 1844)**

Brassica oleracea L. var. *capitata* – Maharashtra (Shaikh et al., 2020)

Brassica rapa L. – Maharashtra (Shaikh et al., 2020)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

Family 2: Reduviidae***Brassivola hystrix* Distant, 1904**

Brassica juncea (L.) Czern. – Delhi (Lakkundi, 1989)

Order: Neuroptera

Neuroptera includes lacewings, mantisflies, antlions and their relatives and both adults and larvae are predatory. They are globally distributed and generalist predators of soft-bodied insects. Recently, Singh et al. (2024a) listed 32 species of aphidophagous neuropterans from India belonging to four families. However, members of only two families, Chrysopidae and Hemerobiidae, were reported to prey on *Lipaphis erysimi* in India. Chrysopidae is the largest family of the order and its members are commonly known as green lacewings. In India, it comprises 24 aphidophagous species preying on 55 species of aphids infesting 84 species of food plants (Singh et al., 2024a). Among them, only four species of lacewings prey on *Lipaphis erysimi* feeding on eight species of food plants in 13 states/union territory. The hemerobiids are commonly known as brown lacewings. In India, two genera, *Hemerobius* and *Micromus*, are recorded as preying on 29 aphid species infesting 48 food plants (Singh et al., 2024a). However, only three species are recorded to prey on *Lipaphis erysimi* infesting five species of plants in India (Table 1). Details on prey and food plant associations are listed below:

Family 1: Chrysopidae***Chrysopa* sp.**

Brassica juncea (L.) Czern. – Tripura (Datta, 2020)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

***Chrysoperla zastrowi sillemi* (Esben – Petersen, 1935)**

Brassica carinata A. Braun – Punjab (Kumar, 2015)

Brassica juncea (L.) Czern. – Assam (Pradhan et al., 2020b); Gujarat (Chakraborty and Korat, 2010); Haryana (Kalra, 1988); Punjab (Soni et al., 2004; Kumar, 2015; Belagalla et al., 2024); Rajasthan (Singh

et al., 2009; Naruka and Ameta, 2015); West Bengal (Ghosh, 1983); Uttarakhand (Verma et al., 2023)

Brassica napus L. – Punjab (Kumar, 2015)

Brassica nigra W.D.J. Koch – Uttar Pradesh (Tiwari et al., 2024b)

Brassica oleracea L. var. *capitata* – Chhattisgarh (Singh, 2014b); Maharashtra (Shaikh et al., 2020); Odisha (Mandal and Patnaik, 2006); Uttar Pradesh (Tiwari et al., 2024b)

Brassica rapa L. – Gujarat (Chaudhary et al., 2022); Maharashtra (Shaikh et al., 2020); Punjab (Kumar, 2015); Rajasthan (Singh et al., 2009); Telangana (Sheileja et al., 2022); Uttar Pradesh (Singh and Kumar, 2000; Tiwari et al., 2024b); Uttarakhand (Joshi and Sharma, 2008)

Eruca vesicaria (L.) Cav. – Punjab (Kumar, 2015)

Raphanus sativus L. – Maharashtra (Shaikh et al., 2020)

***Mallada desjardinsi* (Navás, 1911)**

Brassica juncea (L.) Czern. – Haryana (Kalra, 1988)

Brassica rapa L. – Gujarat (Nehare et al., 2004); Rajasthan (Bakthavatsalam et al., 1994)

***Tumeochrysa indica* Needham, 1909**

Brassica oleracea L. var. *capitata* – West Bengal (Dey and De, 2018)

Raphanus sativus L. – West Bengal (Dey and De, 2018)

Family 1: Hemerobiidae***Hemerobius indicus* Kimmins, 1938**

Brassica rapa L. – West Bengal (Dey et al., 2016)

***Micromus igorotus* Banks, 1920**

Brassica rapa L. – Uttar Pradesh (Tiwari et al., 2024b)

***Micromus timidus* Hagen, 1853 –**

Brassica juncea (L.) Czern. – Manipur (Chitra Devi et al., 2002); Tripura (Datta, 2020); West Bengal (Ghosh, 1983)

Brassica nigra W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Brassica oleracea L. – Manipur (Singh et al., 2002)

Brassica rapa L. – Uttar Pradesh (Tiwari et al., 2024b)

Raphanus sp. – Uttarakhand (Dey, 2015)

Checklist of predators of *Lipaphis pseudobrassicacae* infesting different food plants in India

Predatory arthropods of *L. pseudobrassicacae* belong to three orders of the class Insecta: Coleoptera (Coccinellidae), Diptera (Syrphidae), and Neuroptera (Chrysopidae). A total of 16 species of predators belonging to different taxa were observed to feed on the turnip aphid on five food plant species distributed only in seven states/union territories of India, mostly in Jammu and Kashmir (Table 2). The greatest number of predators belong to the families Syrphidae (10 species) followed by ladybird beetles (5 species) (Table 1). Most of the tritrophic associations (triplets, predators–preys–host plants) of these predators are reported from Jammu and Kashmir.

Class: Insecta**Order: Coleoptera; Family: Coccinellidae*****Cheilomenes sexmaculata* (Fabricius, 1781)**

Brassica juncea (L.) Czern – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *botrytis* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *capitata* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica rapa subsp. *rapa* L. – Jammu and Kashmir (Khan and Shah, 2017)

Raphanus sativus L. – Jammu and Kashmir (Khan and Shah, 2017)

***Coccinella septempunctata* Linnaeus, 1758**

Brassica juncea (L.) Czern – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *botrytis* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *capitata* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica rapa subsp. *rapa* L. – Jammu and Kashmir (Khan and Shah, 2017)

Raphanus sativus L. – Jammu and Kashmir (Khan and Shah, 2017)

***Coccinella transversalis* Fabricius, 1781**

Brassica juncea (L.) Czern. – Tripura (Ghosh et al., 2017)

***Coccinella undecimpunctata* Linnaeus, 1758**

Brassica juncea (L.) Czern. – Bihar (Rahman, 1940)

Brassica juncea (L.) Czern. – Punjab (Rahman, 1940)

***Propylea luteopustulata* (Mulsant, 1850)**

Brassica juncea (L.) Czern – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *botrytis* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *capitata* – Jammu and Kashmir (Khan and Shah, 2017)

Brassica rapa subsp. *rapa* L. – Jammu and Kashmir (Khan and Shah, 2017)

Raphanus sativus L. – Jammu and Kashmir (Khan and Shah, 2017)

Order: Diptera; Family: Syrphidae***Betasyrphus isaaci* (Bhatia, 1933)**

Brassica juncea (L.) Czern. – Bihar (Rahman, 1940)

***Betasyrphus* sp.**

Brassica juncea (L.) Czern. – Karnataka (Chinnu et al., 2023)

***Episyrphus balteatus* (De Geer, 1776)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Episyrphus viridaureus* (Wiedemann, 1824)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1989)

***Eupeodes bucculatus* (Rondani, 1857)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Eupeodes confrater* (Wiedemann, 1830)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Ischiodon scutellaris* (Fabricius, 1805)**

Brassica juncea (L.) Czern. – Delhi (Lal and Gupta, 1953)

***Ischiodon scutellaris* (Fabricius, 1805)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Scaeva latimaculata* (Brunetti, 1923)**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

***Sphaerophoria indiana* Bigot, 1884**

Brassica juncea (L.) Czern. – Punjab (Kumar et al., 1988)

Order: Neuroptera; Family: Chrysopidae***Chrysoperla zastrowi sillemi* (Esben – Petersen, 1935)**

Brassica juncea (L.) Czern – Jammu and Kashmir (Khan and Shah, 2017)

Checklist of parasitoids of *Lipaphis erysimi* infesting different food plants in India

The parasitoids of *Lipaphis erysimi* belong to two families of the order Hymenoptera, Aphelinidae (Superfamily Calcidoidea) and Braconidae (Superfamily Ichneumonoidea) (Tiwari et al., 2024a). A total of 22 species of parasitoids belonging to both families are recorded as parasitizing mustard aphid on 21 host plant species in 25 states/union territories of India. Aphelinidae is represented by 5 species parasitizing the mustard aphids in 7 states/union territory and Braconidae by 17 species parasitizing on 20 host plants in 24 states/union territories of India. These parasitoids were observed to parasitize the turnip aphid on 21 food plant species distributed in 25 states/union territories of India. Most of the tritrophic associations (triplets, parasitoids–hosts–host plants) of these parasitoids are reported from Manipur (15 triplets) followed by Uttar Pradesh (11 triplets) and less than 10 triplets in other states/union territories (Table 3). Detail checklist is given below:

Family 1: Aphelinidae***Aphelinus albipodus* Hayat and Fatima, 1992**

Brassica sp. – Andhra Pradesh (Hayat, 1998)

***Aphelinus asychis* Walker, 1839**

Brassica sp. – Assam (Hayat, 1998); Kerala (Hayat, 1998); Meghalaya (Hayat, 1998); Uttar Pradesh (Hayat, 1998)

***Aphelinus gossypii* Timberlake, 1924**

Trifolium alexandrinum L. – Uttar Pradesh (Ahmad and Singh, 1993)

***Aphelinus* nr. *kurdjumovi* Mercet, 1930**

Brassica sp. – Odisha (Patro, 2007)

***Aphelinus* sp.**

Brassica juncea (L.) Czern. – West Bengal (Ghosh, 1983)

Brassica nigra (L.) W.D.J. Koch – West Bengal (Ghosh et al., 1981)

Table 2: Number of species of predators belonging to different taxa preying on *Lipaphis pseudobrassicae* infesting different number of host plants distributed in number of states/union territories of India.

Order	Families of predators	Number of predator species	Number of host plant species	Number of predator–prey–host plant associations	Number of states/union territories
Coleoptera	Coccinellidae	5	5	18	4
Diptera	Syrphidae	10	1	10	4
Neuroptera	Chrysopidae	1	1	1	1
Total		16	5	29	7

Table 3: Number of species of parasitoids belonging to different taxa parasitizing *Lipaphis erysimi* and *Lipaphis pseudobrassicae* infesting different number of host plants distributed in India.

Families of Parasitoids	Number of parasitoid species	Number of host plant species	Number of parasitoid–host–food plant associations	Number of states/union territories
<i>Lipaphis erysimi</i>				
1. Aphelinidae	5	4	6	7
2. Braconidae	17	21	50	24
<i>Lipaphis pseudobrassicae</i>				
1. Braconidae	3	2	4	3
Total	22	21	60	25

Family 2: Braconidae, subfamily: Aphidiinae

***Aphidius colemani* Viereck, 1912**

Brassica oleracea L. var. *acephala* – Uttar Pradesh (Rafi et al., 2010)

Brassica oleracea L. var. *botrytis* – Bihar (Ahmad and Singh, 2007); Manipur (Subhrani et al., 2010)

Brassica oleracea L. var. *capiatata* – Manipur (Subhrani et al., 2006)

***Aphidius erysimi* (Stary, 1960)**

Brassica sp. – Mizoram (Sarkar, 1991)

Emilia sonchifolia L. (DC.) – Sikkim (Tamili, 1988); West Bengal (Tamili, 1988)

***Aphidius gifuensis* Ashmead, 1906**

Brassica spp. – Odisha (Patro, 2007)

***Aphidius hortensis* Marshall, 1896**

Brassica juncea (L.) Czern. – Manipur (Subhrani et al., 2006)

***Aphidius matricariae* (Haliday, 1834)**

Brassica juncea (L.) Czern. – Tripura (Datta, 2020)

Brassica napus L. – Meghalaya (Stary and Ghosh, 1978; Raychaudhuri et al., 1982); Mizoram (Sarkar, 1991)

Brassica oleracea L. var. *botrytis* – Bihar (Prakash and Rani, 2015)

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996; Devjani and Singh, 1998)

Brassica rapa L. – Uttarakhand (Bisht et al., 2001)

Solanum melongena L. – West Bengal (Tamili, 1988)

***Aphidius* sp.**

Brassica oleracea L. var. *capitata* – Chhattisgarh (Singh, 2014b)

Eruca vesicaria (L.) Cav. – Punjab (Bakhetia and Sharma, 1979)

***Binodoxys brevicornis* (Haliday, 1833)**

Brassica spp. – Odisha (Patro, 2007)

Unknown plant – Karnataka (Stary and Ghosh, 1983)

***Binodoxys eutrichosiphi* (Stary, 1975)**

Brassica rapa L. – Uttar Pradesh (Ahmad and Singh, 1995)

***Binodoxys indicus* (Subba Rao and Sharma, 1958)**

Brassica napus L. – Sikkim (Tamili, 1988)

Brassica rapa L. – Uttar Pradesh (Ahmad and Singh, 1995)

Brassica sp. – Karnataka (Rao et al., 1969)

Calotropis procera (Aiton) Dryand. – Uttar Pradesh (Ahmad and Singh, 1993)

Emilia sonchifolia L. (DC.) – West Bengal (Tamili, 1988)

***Diaeretiella rapae* (McIntosh, 1855)**

Beta vulgaris L. – West Bengal (Tamili, 1988)

Brassica carinata A. Braun – Punjab (Kumar, 2015)

Brassica juncea (L.) Czern – Assam (Pradhan et al., 2020b); Chhattisgarh (Kashyap et al., 2018); Delhi (Akhtar et al., 2010); Gujarat (Kulkarni and Patel, 2001; Patel et al., 2019); Haryana (Singh and Singh, 1994); Himachal Pradesh (Soni et al., 2021); Madhya Pradesh (Lal et al., 2018); Manipur (Subhrani et al., 2006; Rajeshwari and Singh, 2022); Meghalaya (Firake et al., 2013); Punjab (Kumar, 2015); Tripura (Datta, 2020); Uttar Pradesh (Halder et al., 2014); Uttarakhand (Verma et al., 2023)

Brassica napus L. – Meghalaya (Stary and Ghosh, 1978); Punjab (Kumar, 2015)

Brassica nigra (L.) W.D.J. Koch – Karnataka (Rao et al., 1969); Manipur (Singh and Singh, 1986); Meghalaya (Stary and Ghosh, 1978); Nagaland (Singh et al., 1991); Punjab (Atwal et al., 1969)

Brassica oleracea L. – Assam (Rao et al., 1970); Manipur (Devjani and Singh, 1998); Sikkim (Agarwala et al., 1980)

Brassica oleracea L. var. *botrytis* – Bihar (Ahmad and Singh, 1996; Prakash and Rani, 2015); Uttar Pradesh (Ahmad and Singh, 1995; Chaudhary and Singh, 2007); Uttarakhand (Akhtar et al., 2011)

Brassica oleracea L. var. *capitata* – Manipur (Subhrani et al., 2006); Bihar (Ahmad and Singh, 1996); Chhattisgarh (Gauraha, 2021); Manipur (Bijaya et al., 1996); Nagaland (Waluniba et al., 2016)

Brassica rapa L. – Bihar (Ahmad and Singh, 1997); Delhi (Akhtar et al., 2011); Gujarat (Kavad and Korat, 2013); Himachal Pradesh (Nayak et al., 1982; Raj and Sharma, 1993); Jammu and Kashmir (Bhagat and Ahmad, 1991; Ahmad and Ahmad, 2013); Madhya Pradesh (Singh and Rawat, 1981; Veda and Shaw, 1988); Maharashtra (Men and Kandalkar, 1997; Biradar and Dhanorkar, 2004); Meghalaya (Firake et al., 2013); Punjab (Atwal et al., 1969, Atwal et al., 1971); Rajasthan (Chandra and Kushwaha, 1987); Sikkim (Agarwala et al., 1980); Uttar Pradesh (Pandey et al., 1985; Akhtar et al., 2006); Uttarakhand (Kundu et al., 1965; Pervez et al., 2020b); West Bengal (Kundu et al., 1965)

Brassica rapa subsp. *rapa* L. – Uttar Pradesh (Rafi et al., 2010)

Brassica spp. – Himachal Pradesh (Saha et al., 1982); Jammu and Kashmir (Saha, 1988); Karnataka (Rao et al., 1969); Manipur (Singh and Singh, 1986); Meghalaya (Ghosh and Raychaudhuri, 1982; Stary and Ghosh, 1975); Mizoram (Sarkar, 1991); Punjab (Atwal et al., 1969); Uttarakhand (Chakrabarti and Debnath, 2009)

Eruca vesicaria (L.) Cav. – Punjab (Kumar, 2015)

Raphanus sativus L. – Uttar Pradesh (Rafi et al., 2010)

Sinapis arvensis L. – Jammu and Kashmir (Rakhshani et al., 2012)

Trifolium alexandrinum L. – Uttar Pradesh (Rakhshani et al., 2012)

Zea mays L. – Jammu and Kashmir (Rakhshani et al., 2012)

***Ephedrus laevicollis* (Thomson, 1895)**

Brassica juncea (L.) Czern. – Manipur (Subhrani et al., 2006)

Brassica nigra (L.) W.D.J. Koch – Manipur (Singh, 1987)

***Ephedrus minor* Stelfox, 1941**

Brassica nigra (L.) W.D.J. Koch – Manipur (Singh et al., 2011)

***Ephedrus plagiator* (Nees, 1811)**

Brassica oleracea L. var. *capitata* – Manipur (Bijaya et al., 1996; Singh et al., 2011)

***Ephedrus* sp.**

Unknown plant – Sikkim (Stary and Ghosh, 1983)

***Lipolexis gracilis* Forster, 1862**

Brassica rapa L. – Punjab (Sharma and Subba Rao, 1964); Uttar Pradesh (Rafi et al., 2010)

Brassica spp. – Odisha (Patro, 2007)

***Lysiphlebus testaceipes* (Cresson, 1880)**

Brassica sp. – Karnataka (Ramaseshiah et al., 1969)

***Praon volucre* (Haliday, 1833)**

Brassica napus L. – West Bengal (Tamili, 1988)

Brassica spp. – Jammu and Kashmir (Stary and Bhagat, 1978); Meghalaya (Samanta, 1986); Mizoram (Sarkar, 1991); Odisha (Patro, 2007); Indet – Himachal Pradesh (Stary and Ghosh, 1983)

Checklist of parasitoids of *Lipaphis pseudobrassicacae* infesting different food plants in India

Only two identified species of Braconidae were observed to parasitize *Lipaphis pseudobrassicacae* in India on two host plant species distributed only in Jammu and Kashmir, Karnataka and Punjab (Table 3) as listed below:

Family: Braconidae

***Aphidius erysimi* (Stary, 1960)**

Brassica juncea (L.) Czern – Jammu and Kashmir (Khan and Shah, 2017)

Brassica oleracea L. var. *capitata* – Jammu and Kashmir (Khan and Shah, 2017)

***Aphidius* sp.**

Brassica oleracea L. var. *capitata* – Karnataka (Chinnu et al., 2023)

***Diaeretiella rapae* (McIntosh, 1855)**

Brassica juncea L. – Punjab (Kumar et al., 1989)

Discussion

There is a large number of arthropods that either feed on *L. erysimi* (predators) or parasitize it (parasitoids) in nature, keeping its population under control. Among them, few species of parasitoids such as *A. matricariae* and *D. rapae* and predators such as *C. septempunctata*, *C. zastrowi sillemi*, *I. scutellaris*, etc., have been used as bioagents in its biological control throughout the world (Singh and Singh, 2016). As natural control agents, all the recorded species of predators and parasitoids, particularly the ladybird beetles, play major roles because of their species diversity and by possessing high foraging performance, predatory efficiency and high reproductive potential (Chellappan and Ranjith, 2023). One hundred species of predators and 22 species of parasitoids of *L. erysimi* and *L. pseudobrassicacae* recorded may regulate the pests on brassica oilseed and vegetable crops, reducing the yield loss in India if they are properly managed in the field population via cultural practices.

Acknowledgements

I would like to extend our gratitude to both the reviewers, Prof. Oldřich Nedvěd, Faculty of Science, University of South Bohemia, Institute of Entomology, Biological Center, Academy of Sciences of the Czech Republic and Dr. J. T. D. Caleb, Department of Anatomy, Saveetha Medical College & Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India for their valuable comments/suggestions that helped me to improve the manuscript. The author also thanks Dr. Erica DeMilio, Department of Zoology, School of Natural Sciences, Ryan Institute, National University of

Ireland Galway, Ireland for her valuable scientific comments and language editing.

Author contributions

The author himself collected all the data from different resources and prepares the manuscript.

Funding

No specific grant was received from any funding agency.

Conflict of interest

The author declares that there are no conflicting issues related to this review article.

References

- Afroze, S. (1999). Preferential predation by *Anegleis cardoni* (Weise) (Coccinellidae: Coleoptera) on some aphid species. *Journal of Aphidology*, 13 (1–2): 95–97.
- Agarwala, B. K., Dutta, S. and Raychaudhuri, D. N. (1982). An account of syrphid (Diptera: Syrphidae) predators of aphids (Homoptera: Aphididae) available in Darjeeling district of West Bengal and Sikkim. *Pranikee*, 3: 15–21.
- Agarwala, B. K., Bhaumik, A. K. and Gilbert, F. S. (1989). Relative development and voracity of six species of aphidophagous syrphids in cruciferous crops. *Proceedings of the Indian Academy of Sciences (Animal Science)*, 98 (4): 267–274. <https://doi.org/10.1007/BF03179408>
- Agarwala, B. K., Das, S. and Senchowdhuri, M. (1988). Biology and food relations of *Micraspis discolor* (F.): an aphidophagous coccinellid in India. *Journal of Aphidology*, 2 (1–2): 7–17.
- Agarwala, B. K., Ghosh, D., Das, S. K., Poddar, S. C. and Raychaudhuri, D. N. (1981). Parasites and predators of aphids (Homoptera: Aphididae) from India. V. New records of two aphidiid parasites, nine arachnids and one dipteran predator from India. *Entomon*, 6 (3): 233–238.
- Agarwala, B. K., Laska, P. and Raychaudhuri, D. N. (1984). Prey records of aphidophagous syrphid flies from India (Diptera: Syrphidae). *Acta Entomologica Bohemoslovaca*, 81 (1): 15–21.
- Agarwala, B. K., Raychaudhuri, D. and Raychaudhuri, D. N. (1980). Parasites and predators of aphids in Sikkim and Manipur (Northeast India). III. *Entomon*, 5 (1): 39–42.
- Agarwala, B. K., Saha, J. L. and Mahapatra, S. K. (1987). Three new species of parasitoids (Hymenoptera: Aphidiidae) from India. *Entomon*, 12 (4): 329–333.
- Ahmad, D. and Bhat, M. R. (1986). Distribution and host range of some aphidophagous syrphid flies in Kashmir. *Geobios New Reports*, 5: 165–166.
- Ahmad, M. E., Kumar, S., Parween, N. and Rakhshan (2012). Screening of aphidophagous coccinellids (Coleoptera: Coccinellidae) from northeast Bihar. *Journal of Advanced Zoology*, 33 (2): 103–112.
- Ahmad, M. E. and Kumari, K. (2024). Prey records of aphidophagous syrphids with their prey and host plants in India: A review. *Journal of Biological Control*, 38 (2): 117–137. <https://doi.org/10.18311/jbc/2024/35964>
- Ahmad, M. E. and Singh, R. (1993). Records of aphid parasitoids from north eastern Uttar Pradesh. *Journal of Aphidology*, 6–7: 7–14.
- Ahmad, M. E. and Singh, R. (1995). Records of Macrosiphini of Northeastern Uttar Pradesh and its relationship with food plants and natural enemies. *Journal of Aphidology*, 9 (1–2): 80–86.
- Ahmad, M. E. and Singh, R. (1996). Records of aphid parasitoids from the North Bihar and associations with their hosts and food plants. *Journal of Advanced Zoology*, 17 (1–2): 26–33.
- Ahmad, M. E. and Singh, R. (1997). Records of aphids and their food plants, parasitoids and hyperparasitoids from north Bihar. *Journal of Advanced Zoology*, 18 (1): 54–61.
- Ahmad, M. E. and Singh, R. (2007). Tritrophic relations of aphids (Homoptera: Aphididae) infesting vegetable plants in north east Uttar Pradesh. *Journal of Aphidology*, 21 (1–2): 89–95.
- Ahmad, M. J. and Ahmad, S. B. (2013). Dynamics of *Diaeretiella rapae* (M'Intosh) (Braconidae: Aphidiidae) and its hyperparasitoid on mustard aphid, *Lipaphis erysimi* (Kalt.) infesting brown mustard, *Brassica campestris* Linn. in Kashmir, India. *Journal of Biological Control*, 27 (4): 247–252. <https://doi.org/10.18311/jbc/2013/3253>
- Akhtar, M. S., Dey, D. and Usmani, M. K. (2010). Spatial distribution of mustard aphid *Lipaphis erysimi* (Kaltenbach) Vis-à-vis its parasitoid, *Diaeretiella rapae* (M'Intosh). *World Applied Sciences Journal*, 11 (3): 284–288.
- Akhtar, M. S., Choudhury, R. A. and Shuja-Uddin. (2006). Natural enemies of mustard aphid, *Lipaphis erysimi* (Kaltenbach) in Aligarh (India). *Environment and Ecology*, 24S: 781–784.
- Akhtar, M. S., Rafi, U., Usmani, M. K. and Dey, D. (2011). A review of aphid parasitoids (Hymenoptera: Braconidae) of Uttar Pradesh and Uttarakhnad, India. *Biology and Medicine*, 3 (2 Special Issue): 320–323.
- Ali, A., Rizvi, P. Q. and Khan, F. R. (2009). On the predation of aphids by *Ischiodon scutellaris* (Diptera: Syrphidae) under natural environment. *Bionotes*, 11 (3): 95–96.

- Atwal, A. S. and Sethi, S. L. (1963). Predation by *Coccinella septempunctata* L. on the cabbage aphid, *Lipaphis erysimi* (Kalt.) in India. *Journal of Animal Ecology*, 32 (3): 481–488.
<https://doi.org/10.2307/2604>
- Atwal, A. S., Chaudhary, J. P. and Ramzan, M. (1969). Some preliminary studies in India on the bionomics and rate of parasitisation of *Diaeretiella rapae* Curtis (Braconidae: Hymenoptera), a parasite of aphids. *Journal of Research, Punjab Agricultural University*, 6 (1): 177–182.
- Atwal, A. S., Chaudhary, J. P. and Ramzan, M. (1971). Mortality factors in the natural population of cabbage aphid, *Lipaphis erysimi* (Kalt.) (Aphididae: Homoptera) in relation to parasites, predators and weather conditions. *Indian Journal of Agricultural Science*, 41 (5): 507–510.
- Bakhietia, D. R. C. and Sekhon, B. S. (1989). Insectpests and their management in rapeseed mustard. *Journal Oilseed Research*, 6 (3): 269–299.
- Bakhietia, D. R. and Sharma, A. K. (1979). Preliminary observations on aphids infestation on *Eruca sativa* Mill. *Indian Journal of Entomology*, 41 (3): 288–289.
- Bakthavatsalam, N. and Varshney, R. (2023). Chrysopids (Neuroptera: Chrysopidae), In: Omkar (Ed.), *Insect Predators in Pest Management*. CRC Press, Taylor and Francis Group, LLC. pp. 161–174.
<https://doi.org/10.1201/9781003370864-6>
- Bakthavatsalam, N., Singh, S. P., Pushpalatha, N. A. and Bhummannavar, B. S. (1994). Life tables of four species of chrysopids (Neuroptera: Chrysopidae). *Journal of Entomological Research*, 18 (4): 357–360.
- Bali, R. (1987). Coccinellids (Coleoptera) of Kashmir valley, distribution and host range. *Geobios new reports*, 6: 187–189.
- Ballal, C. R., Gupta, T. and Varshney, R. (2023). The flower bugs (Anthocoridae), In: Omkar (Ed.), *Insect Predators in Pest Management*. CRC Press, Taylor and Francis Group, LLC. pp. 73–106.
<https://doi.org/10.1201/9781003370864-3>
- Bana, J. K., Jat, B. L. and Bajya, D. R. (2012). Seasonal incidence of major pests of cabbage and their natural enemies in relation to weather parameters. *Indian Journal of Entomology*, 74 (3): 236–240.
- Belagalla, N., Kaur, R., Sangha, K. S. and Kang, B. K. (2024). Predating response of *Chrysoperla zastrowi sillemi* (Esben-Peterson) on raya aphid *Lipaphis erysimi* (Kaltenbach). *Journal of Advanced Zoology*, 45 (3): 623–630.
<https://doi.org/10.56557/upjz/2024/v45i63957>
- Bhagat, R. C. and Ahmad, M. N. (1991). Aphidiid parasitoids (Hymenoptera) of aphids (Homoptera) of Jammu – new records, host range and biological notes. *Journal of Aphidology*, 5 (1–2): 90–96.
- Bhagat, R. C. and Lone, M. A. (1984). New records and host range of predators of aphids (Aphididae: Homoptera) in Kashmir valley India. *Science and Culture*, 50 (4): 364–366.
- Bhagat, R. C. and Matta, T. A. (2002). Host range and diversity of aphidophagous insect (predators) of Kashmir Himalaya, with new host aphid/plant records, In: Khan, M. A. (Ed.), *Environment Biodiversity and Conservation*. A. P. H. Publishing Corporation New Delhi. pp. 269–282.
- Bhat, D. M. and Bhagat, R. C. (2017). Host range and diversity of syrphid predators (Insecta: Diptera) of aphids on vegetable crops of Kashmir, with new host aphid/ plant records. *Trends in Biosciences*, 10 (6): 1446–1448.
- Bhat, D. M. (2008). Studies on insect parasites and predators of some insect pests of vegetable crops of Kashmir Valley. Ph.D. thesis, P. G. Department of Zoology, University of Kashmir, Srinagar, J&K (India). 193 pp.
<http://hdl.handle.net/10603/72857>
- Bhat, D. M. (2017). Host range and diversity of coccinellid (Coleoptera) predators of aphid pests in vegetable crop ecosystems of Kashmir, with new host aphid/plant records. *Journal of Entomological Research*, 41 (2): 183–186.
<https://doi.org/10.5958/09744576.2017.00029.9>
- Bijaya, P. D., Devi, C., Varatharajan, R. and Singh, T. K. (1996). On the density and composition of natural enemies of cabbage aphids from Manipur. *Journal of Advanced Zoology*, 17 (2): 74–78.
- Biradar, V. K. and Dhanorkar, B. K. (2004). Seasonal dynamics of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) and its parasitoid, *Diaeretiella rapae* (Mcintosh) (Hymenoptera: Braconidae) in Marathwada region of Maharashtra. *Journal of Aphidology*, 18 (1–2): 1–4.
- Bisht, R. S., Rana, D. S. and Mishra, P. N. (2001). Bio-Ecology of predators and parasitoids of aphids in Garhwal Hills. *Journal of Aphidology*, 15 (1–2): 123–128.
- Bisht, R. S., Sharma, R. K. and Dev, P. (2006). Vertical distribution and activity of aphidophagous syrphids (Diptera: Syrphidae) in Garhwal Himalayas. *Journal of Aphidology*, 20 (1–2): 25–29.
- Boopathi, T., Singh, S. B., Dutta, S. K., Dayal, V., Singh, A. R., Chowdhury, S., Ramakrishna, Y., Shakuntala, I. and Lalhruaipuii, K. (2020). Biology, predatory potential, life table, and field evaluation of *Propylea dissecta* (Coleoptera: Coccinellidae), against *Lipaphis erysimi* (Homoptera: Aphididae) on broccoli. *Journal of Economic Entomology*, 113 (1): 88–97.
<https://doi.org/10.1093/jee/toz272>
- Borah, R. and Dutta, S. K. (2010). Feeding potential of *Coccinella transversalis* F. on different morphs of mustard aphid, *Lipaphis erysimi* (Kaltenbach). *Journal of Biological Control*, 24 (3): 271–273.

- Borah, R. and Dutta, S. K. (2010). Predatory potential of *Episyrphus balteatus* (De Geer), an effective predator of *Lipaphis erysimi* (Kaltenbach). *Journal of Biological Control*, 24 (2): 173–174.
- Capinera, J. L. (2001). *Handbook of Vegetable Pests*. Gulf Professional Publishing. pp. 327–328.
- Chakrabarti, S. and Debnath, M. (2009). Diversity of aphidophagous parasitoids (Insecta) of northwest and western Himalayas, India, In: Hartmann, M. and Weipert, J. (Eds.), *Biodiversität und Naturschutz im Himalaya, III*. Naturkundemuseum Erfurt. pp. 441–454.
- Chakrabarti, S., Sarkar, S. and Debnath, M. (2012). Diversity, bioecology and biosystematics of aphidophagous predators of eastern Himalaya and northeast India, In: Hartmann, M. and Weipert, J. (Eds.), *Biodiversität und Naturschutz im Himalaya, IV*. Naturkundemuseum Erfurt. pp. 129–147.
- Chakraborty, D. and Korat, D. M. (2010). Feeding efficiency of green lacewing, *Chrysoperla carnea* (Stephens) on different species of aphids. *Karnataka Journal of Agricultural Sciences*, 2 (5): 793–794.
- Chakraborty, D. (2012). Biodiversity, life-history and biotic potential of predatory coccinellids of Anand region of Gujarat. Ph.D. thesis, Anand Agricultural University, Anand, Gujarat. 204 pp.
- Chand, A., Khinchi, S. K., Kumawat, K. C., Hussain, A. and Sharma, S. L. (2022). Quantitative and qualitative status of insect pests of mustard, *Brassica juncea* (L.) Czern and Coss and their natural enemies. *The Pharma Innovation Journal*, SP-11 (1): 412–415.
- Chandra, S. and Kushwaha, K. S. (1987). Impact of environmental resistance on aphid complex of cruciferous crops under the agroclimatic conditions of Udaipur II. Biotic components. *Indian Journal of Entomology*, 49 (1): 86–113.
- Chaudhari, P., Dabhi, M. R. Patel, N. B. and Raghunandan, B. L. (2022). Predatory efficiency of *Chrysoperla zastrowi sillemi* (Esben-Peterson) on different hosts. *Frontiers in Crop Improvement*, 10: 307–309.
- Chaudhary, H. C. and Singh, R. (2007). Tritrophic relations of aphids (Homoptera: Aphididae) infesting vegetable plants in northeast Uttar Pradesh. *Journal of Aphidology*, 21 (1–2): 89–95.
- Chaudhary, H. C. and Singh, R. (2012). Records of the predators of aphids (Homoptera: Aphididae) in eastern Uttar Pradesh. *Journal of Aphidology*, 25 and 26: 13–30.
- Chellappan, M. and Ranjith, M. T. (2023). Insect predators, In: Omkar (Ed.), *Insect predators in pest management*. CRC Press, Taylor and Francis Group, LLC. pp. 1–54. <https://doi.org/10.1201/9781003370864-1>
- Chinnu, V. S. Mulimani, V., Shanas, S. and Sumithramma, N. (2023). Exploration of natural enemy fauna of aphids and associated ant species from eastern dry zone of Karnataka, India. *Entomon*, 48 (4): 545–552. <https://doi.org/10.33307/entomon.v48i4.990>
- Chitra Devi, L., Singh, T. K. and Varatharajan, R. (1997). Impact of predators on the field population of *Lipaphis erysimi* (Kalt.) infesting mustard. *Journal of Aphidology*, 11 (1): 107–110.
- Chitra Devi, L., Singh, T. K. and Varatharajan, R. (2002). Role of natural enemies in the management of *Lipaphis erysimi* (Kalt.) on *Brassica juncea* var. *rugosa* (Linn.). *Journal of Biological Control*, 16 (1): 27–30.
- Chitra Devi, L., Varatharajan, R., Singh, T. K. (1996). Prey-predator relations of *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) and aphidophagous syrphids on mustard. *Uttar Pradesh Journal of Zoology*, 16 (1): 9–12.
- Choudhary, S., Meena, R. K., Jat, B. L., Hussain, A. and Sharma, P. (2023). Population dynamics of mustard aphid, *Lipaphis erysimi* (Kaltenbach) on cauliflower in relation to biotic and abiotic factors. *International Journal of Environment and Climate Change*, 13 (11): 2322–2328. <https://doi.org/10.9734/ijec/2023/v13i113395>
- Chouhan, K. S., Patel, B. H., Korat, D. M. and Chakraborty, D. (2011). Predatory potential of syrphid fly, *Ischiodon scutellaris* (Fab.) on four different species of aphids in laboratory conditions. *Insect Environment*, 17 (3): 107–108.
- Damitre, L. (2019). Effect of different temperatures and pesticides on the biological attributes of major coccinellid beetles predating on mustard aphid *Lipaphis erysimi* Kaltenbach. Ph.D. thesis, Nagaland University, Medziphema, Nagaland. 78 pp. <http://hdl.handle.net/10603/315226>
- Das, R. (2020). A field study on insect pest complex of brassicaceous crops in some areas of Cachar, Assam. *Journal of Entomology and Zoology Studies*, 8 (4): 2043–2045.
- Datta, N. (2020). The mustard aphid *Lipaphis erysimi* (Kaltenbach) and its natural enemies in Tripura: a review. *Research Journal of Life Sciences, Bioinformatics, Pharmaceuticals and Chemical Sciences*, 6 (5): 17–28. <https://doi.org/10.26479/2020.0605.02>
- Debnath, N. (1991). Studies on some aphidophagous insects in northwest and western Himalays. Ph.D. thesis, University of Kalyani, India. <http://hdl.handle.net/10603/211339>
- Devi, Y. R., Kalita, J. and Singh, T. K. (2011). Biological control potential of an aphidophagous syrphid, *Episyrphus balteatus* De-Geer (Diptera: Syrphidae) on mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) on cabbage ecosystem in Manipur. *Journal of Experimental Sciences*, 2 (12): 13–16.
- Devjani, P. and Singh, T. K. (1998). Ecological succession of aphids and their natural enemies on cauliflower in Manipur. *Journal of Aphidology*, 12 (1–2): 45–51.
- Devjani, P. and Singh, T. K. (2006). Larval development and prey consumption of five aphidophagous predators in relation to two aphid species, In: *Advances in Indian Entomology: Productivity and Health a Silver Jubilee*, Supplement No. 3, Volume-II, Insect Environment. pp. 121–125.

- Dey, S. R. (2015). Seasonal occurrence and altitudinal distribution of Neuroptera (Insecta) in Uttarakhand, India. *The Beats of Natural Sciences*, 2 (4): 1–13.
- Dey, S. R. and De, M. (2018). Report on winter crop plant–aphid–aphidophagous Chrysopidae (Neuroptera: Insecta) association from Murshidabad district, West Bengal, India. *International Journal of Experimental Research and Review*, 16: 7–13.
<https://doi.org/10.52756/ijerr.2018.v16.002>
- Dey, S. R., Sikdar, S. R., Ghose, T. K. and De, M. (2016). Study of the efficiency of neuropteran predator, *Hemerobius indicus* Kimmins as potential biocontrol agent of notorious aphid, *Lipaphis erysimi* (Kaltenbuch) on *Brassica campestris* Linn. (cv. B–9). *International Journal of Experimental Research and Review*, 7: 21–31.
- Dwivedi, S. A., Nameirakpam, L. and Tomer, A. (2021). Brassica–aphid interaction: modulated challenges and sustainable approach for management, In: Aminul Islam, A. K. M., Hossain, M. A. and Mominul Islam, A. K. M. (Eds.), *Brassica Breeding and Biotechnology*. IntechOpen. pp. 1–20.
<https://doi.org/10.5772/intechopen.96903>
- Favret, C. (2024). Aphid Species File. <https://Aphid.SpeciesFile.org> (Accessed 5 August 2024).
- Firake, D. M., Behere, G. T. and Thakur, N. S. A. (2013). Eco-Friendly management of mustard aphid, *Lipaphis erysimi*: a most destructive pest of brassicaceous crops. *Popular Kheti*, 1 (3): 31–33.
- Gangopadhyay, R. N. (1980). Studies on the aphids of Tripura with special reference to biology and control of some of the economically important species. Ph.D. thesis, Calcutta University, Kolkata, West Bengal, India. 240 pp.
<http://hdl.handle.net/10603/162502>
- Gauraha, R. (2021). Studies on insect pests and associated defenders on cabbage, *Brassica oleracea* var. capitata at Raipur, Chhattisgarh. Ph.D. thesis, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh. 186 pp.
<https://krishikosh.egranth.ac.in/handle/1/5810180214>
- Gautam, R. D. (1990). Mass multiplication technique of coccinellid predator, ladybird beetle (*Brumoides suturalis*). *Indian Journal of Agricultural Sciences*, 60 (11): 747–750.
- GBIF (2024). The Global Biodiversity Information Facility. <https://www.gbif.org> (Accessed 5 August 2024).
- Ghosh, A. K. and Raychaudhuri, D. N. (1982). Ecology of natural enemy complex of Aphidoidea (Homoptera) in some areas of eastern India. *Proceedings of the Symposium on Ecology of Animal Population, Zoological Survey of India*, 3: 55–69.
- Ghosh, D. (1983). Studies on mustard aphid *Lipaphis erysimi* (Kltb.) (Homoptera: Insecta) and its natural enemy complex in some parts of West Bengal. Ph.D. thesis, Calcutta University, Kolkata, West Bengal, India. 207 pp.
<http://hdl.handle.net/10603/163460>
- Ghosh, D., Debnath, N. and Chakrabarti, S. (1991). Predators and parasites of aphids from northwest and western Himalaya III. Twenty-five species of coccinellidae (Coleoptera: Insecta) from Garhwal and Kumaon ranges. *Records of Zoological Survey of India*, 88 (2): 177–188.
<https://doi.org/10.26515/rzsi/v88/i2/1991/161342>
- Ghosh, D., Poddar, S. C. and Raychaudhuri, D. N. (1981). Natural enemy complex of *Aphis craccivora* Koch and *Lipaphis erysimi* (Kalt.) in and around Calcutta. *Science and Culture*, 47 (1): 58–60.
- Ghosh, S., Behere, G. T. and Agarwala, B. K. (2017). Molecular characterization of ladybird predators (Coleoptera: Coccinellidae) of aphid pests (Homoptera: Aphididae) in North East India. *Current Science*, 113 (9): 1755–1759.
<https://doi.org/10.18520/cs/v113/i09/1755-1759>
- Goswami, T. N., Anil, C. N. and Chandran, N. (2016). Ladybird beetles in major rabi oil seeds and pulse crops at Sabour, Bihar. *International Journal of Science, Environment and Technology*, 5 (4): 2382–2386.
- Gour, I. S. and Pareek, B. L. (2003). Biology and predatory potential of *Coccinella septempunctata* Lin. on mustard aphid in the semi-arid region of Rajasthan. *Annals of Biology*, 19 (2): 225–229.
- Gurung, B., Ponnusamy N. and Pal, S. (2019). Species diversity of predaceous coccinellids in different crop ecosystems under the hilly and terai region of West Bengal (India). *Ecology, Environment and Conservation*, 25 (2): 636–642.
- Halder, J., Rai, A. B. and Kodandarm, M. H. (2014). Parasitization preference of *Diaeretiella rapae* (Hymenoptera: Braconidae) among different aphids in vegetable ecosystem. *Indian Journal of Agricultural Sciences*, 84 (11): 1431–1433.
<https://doi.org/10.56093/ijas.v84i11.44663>
- Hayat, M. (1998) Aphelinidae of India (Hymenoptera: Chalcidoidea): a taxonomic revision. *Memoirs on Entomology International*, 13: 1–416.
- Heimpel, G. E. and Mills, N. J. (2017). *Biological Control: Ecology and Applications*. Cambridge University Press, UK. 386 pp.
<https://doi.org/10.1017/9781139029117>
- Hugar, P. G., Anandhi, P., Varma, S. and Sarvanan, L. (2008). Seasonal incidence of important insect pests of mustard and their natural enemies in relation to weather parameters, in Allahabad region. *Journal of Entomological Research*, 32 (2): 109–112.
- Jadhav, D. S. and Shukla, A. (2015). Feeding potential of *Coccinella transversalis* (Fabricius) on aphids. *Indian Journal of Entomology*, 77 (2): 197–198.
<https://doi.org/10.5958/09748172.2015.00039.5>
- Joshi, P. C and Sharma, P. K. (2008). First records of coccinellid beetles (Coccinellidae) from the Haridwar, (Uttarakhand), India. *The Natural History Journal of Chulalongkorn University*, 8 (2): 157–167.

- Joshi, S. and Ballal, C. R. (2013). Syrphid predators for biological control of aphids. *Journal of Biological Control*, 27 (3): 151–170.
- Joshi, S., Ballal, C. R. and Rao, N. S. (1999a). Biotic potential of three coccinellid predators on six different aphid hosts. *Journal of Entomological Research*, 23 (1): 1–7.
- Joshi, S., Ballal, C. R. and Rao, N. S. (1999b). Evaluation of biotic potential of syrphid predators *Ischiodon scutellaris* (Fabricius) and *Paragus serratus* (Fabricius) (Diptera: Syrphidae). *Journal of Aphidology*, 13 (1–2): 9–16.
- Joshi, S., David, K. J. and Sachin, K. (2023). Syrphid predators (Diptera: Brachycera), In: Omkar (Ed.), *Insect Predators in Pest Management*. CRC Press, Taylor and Francis Group, LLC. pp. 229–244. <https://doi.org/10.1201/9781003370864-9>
- Kale, P., Bisen, A., Naikwadi, B., Bhure, K. and Undirwade, D. B. (2020). Diversity study of aphids and associated predatory fauna occurred in major kharif and rabi crop ecosystems of Akola, Maharashtra, India. *International Journal of Chemical Studies*, 8 (4): 3868–3876. <https://doi.org/10.22271/chemi.2020.v8.i4aw.10254>
- Kalra, V. K. (1988). Population dynamics of various predators associated with mustard aphid, *Lipaphis erysimi* (Kalt.). *Journal of Biological Control*, 2 (2): 77–79.
- Karnatak, A. K. and Thorat, P. V. (2006). Feeding potential of a ladybird, *Coccinella septempunctata* and a syrphid, *Xanthogramma scutellare* on mustard aphid, *Lipaphis erysimi*. *Journal of Applied Bioscience*, 32 (1): 123–125.
- Kashyap, N., Painkra, G. P., Painkra, K. L. and Bhagat, P. K. (2018). Insect–pests succession, natural enemies and their correlation with weather parameters in mustard crop. *Journal of Plant Development Sciences*, 10 (10): 563–568.
- Kavad, N. K. and Korat, D. M. (2013). Activity of *Diaeretiella rapae* (McIntosh) – Effect of biotic and abiotic factors. *Insect Environment*, 19 (1): 69–72.
- Khan, A. A. and Shah, M. A. (2017). Records of aphid and their natural enemies in agro–ecosystem with special reference to horticultural ecosystem of Kashmir. *Journal of Entomology and Zoology Studies*, 5 (4): 189–203.
- Khan, A. A., Zaki, F. A., Khan, Z. H. and Mir, M. A. (2009). Biodiversity of predaceous beetles (Coleoptera: Coccinellidae) in Kashmir. *Journal of Biological Control*, 23 (1): 43–47.
- Kishor, R., Malik, Y. P., Sirajuddin, Mishra, P. K., Kushwaha, D. and Verma, S. K. (2023). Correlation between the seasonal incidence of aphids and coccinellids on the Indian mustard in different varieties and sowing dates. *International Journal of Environment and Climate Change*, 13 (11): 1452–1466. <https://doi.org/10.9734/ijecce/2023/v13i113297>
- Kolhekar, S., Paikra, M., Gupta, K. and Ganguli, J. (2019). Feeding potential of coccinellid predator, *Menochilus sexmaculatus* (Fabricius) (Coleoptera: Coccinellidae) on mustard aphid, *Lipaphis erysimi*. *Journal of Pharmacognosy and Phytochemistry*, SP2: 35–36.
- Kulkarni, A. V. and Patel, I. S. (2001). Seasonal abundance of mustard aphid (*Lipaphis erysimi*) and associated biogents in Indian mustard (*Brassica juncea*) crop. *The Indian Journal of Agricultural Sciences*, 71 (10): 681–682.
- Kumar, A., Kapoor, V. C. and Lriska, P. (1987). Immature stages of some aphidophagous syrphidflies of India (Insecta, Diptera, Syrphidae). *Zoologica Scripta*, 16 (1): 83–88. <https://doi.org/10.1111/j.1463-6409.1987.tb00055.x>
- Kumar, A., Kapoor, V. C. and Mahal, M. S. (1988). Population buildup and dispersion of immature stages of aphidophagous syrphids (Syrphidae: Diptera) on raya, *Brassica juncea* Coss. *Journal of Insect Science*, 1: 39–84.
- Kumar, A., Kapoor, V. C. and Mahal, M. S. (1989). Pupal population of aphidophagous syrphids and their hyperparasitism on *Brassica juncea* Coss. *Journal of Insect Science*, 2: 114–118.
- Kumar, S. (2015). Relative abundance of turnip aphid and the associated natural enemies on oilseed *Brassica* genotypes. *Journal of Agricultural Science and Technology*, 17: 1209–1222.
- Kumar, S., Ahmad, M. E. and Rakhshan, R. (2016). Influence of prey species on feeding preference, post–embryonic development and life span of *Cheilomenes sexmaculata* (Fabricius). *European Scientific Journal*, 12 (36): 403–410. <https://doi.org/10.19044/esj.2016.v12n36p403>
- Kundu, G. G., Sharma, V. K., Anand, R. K., Rai, S. (1965). New record of *Diaeretiella rapae* as a parasite of mustard aphid *Lipaphis erysimi* (Hemiptera: Aphididae). *Indian Journal of Entomology*, 27 (4): 497–498.
- Lakhanpal, G. C. and Raj, D. (1998). Predation potential of coccinellid and syrphid on important aphid species infesting rapeseed in Himachal Pradesh. *Journal of Entomological Research*, 22 (2): 181–190.
- Lakkundi, N. H. (1989). Assessment of reduviids for their predation and possibilities of their utilization in biological control. Ph.D. thesis, IARI, Division of Entomology, New Delhi, India.
- Lal, B., Nayak, M. K., Srivastava, A. K. and Tomar, D. S. (2018). Seasonal incidence of major insect pests of mustard and their natural enemies. *International Journal of Chemical Studies*, 6 (3): 3658–3661.
- Lal, R. and Gupta, S. B. L. (1953). Morphology of the immature Stages of *Sphaerophoria scutellaris* (Fabr.) (Syrphidae: Diptera) with notes on its biology. *Indian Journal of Entomology*, 13 (3): 207–218.

- Maji, A., Pal, S., Gurung, B., Hatterjee, M. C. and Sahoo, S. K. (2022). Diversity of aphids and their predatory coccinellids from West Bengal. *Indian Journal of Entomology*, 85 (2): 332–336.
<https://doi.org/10.55446/IJE.2022.682>
- Mandal, S. M. A. and Patnaik, N. C. (2006). Insect pests and natural enemies associated with cabbage in the coastal belt of Orissa. *Journal of Plant Protection and Environment*, 3 (1): 76–80.
- Manpoong, N. S., Firake, D. M., Behere, G. T. and Rajesh, T. (2016). Biological attributes and feeding potential of three dominant predators of *Lipaphis erysimi* (Kaltenbach). *Journal of Biological Control*, 30 (3): 190–194.
<https://doi.org/10.18311/jbc/2016/15601>
- Men, U. B. and Kandalkar, H. G. (1997). Records on *Syrphophagous aphidivorous* Mayr (Hymenoptera: Encyrtidae), a hyperparasitoid of a primary parasitoid, *Diaeretiella rapae* (M'Intosh) (Hymenoptera: Braconidae) on mustard aphid, *Lipaphis erysimi* (Kaltenbach) (Homoptera: Aphididae). *PKV Research Journal*, 21: 197–198.
- Mir, R. A. (2013). Composition biology and predation of ladybird beetles on aphids in agro-ecosystem of Kashmir Ph.D. thesis, Aligarh Muslim University, Aligarh, Uttar Pradesh, India. 118 pp.
<http://hdl.handle.net/10603/63173>
- Mishra, M. and Gaurav, K. (2022). Management of mustard aphid (*Lipaphis erysimi* (Kalt.)) by manipulating sowing dates. *Journal of Pharmaceutical Negative Results*, Special Issue, 13 (9): 8489–8498.
- Mishra, M. and Mukharjee, U. (2020). Effect of environmental factors on natural enemies of mustard aphid. *Journal of Entomology and Zoology Studies*, 8 (5): 346–349.
- Mishra, S. K. and Kanwat, P. M. (2017). Biology and predatory potential of *Coccinella septempunctata* Linn. on *Lipaphis erysimi* Kalt. under controlled conditions. *Plant Archives*, 17 (2): 1099–1104.
- Mishra, S. K. and Kanwat, P. M. (2018). Seasonal incidence of mustard aphid, *Lipaphis erysimi* (Kalt) and its major predator on mustard and their correlation with abiotic factors. *Journal of Entomology and Zoology Studies*, 6 (3): 831–836.
- Naruka, P. and Ameta, O. P. (2015). Biology of *Chrysoperla zastrowi arabica* (Henery et al.) on different prey. *Indian Journal of Applied Entomology*, 29 (1): 61–65.
- Nayak, M. R. C., Basu, M. and Raychaudhuri, D. N. (1982). Parasites and predators of aphids (Homoptera: Aphididae) from India–V. Seven coleopteran and four dipteran predators from Orissa. *Pranikee*, 3: 7–14.
- Nehare, S. K., Deotale, V. Y., Deotale, R. O. and Dawane, P. N. (2004). Biology and predatory potential of *Mallada boninensis* (Okamoto) against sucking pests. *Journal of Soil and Crops*, 14 (2): 427–432.
- Nyffeler, M. and Birkhofer, K. (2017). An estimated 400–800 million tons of prey are annually killed by the global spider community. *The Science of Nature*, 104 (3–4): 30, pp. 12.
<https://doi.org/10.1007/s00114-017-1440-1>
- Omkar and Bind, R. B. (1995). Records of aphids–natural enemies complex of Uttar Pradesh. IV. The coccinellids. *Journal of Advanced Zoology*, 16 (2): 67–71.
- Omkar and Mishra, G. (2005). Preference–performance of a generalist predatory ladybird: a laboratory study. *Biological Control*, 34 (2): 187–195.
<https://doi.org/10.1016/j.biocontrol.2005.05.007>
- Omkar and Pervez, A. (1999). New record of coccinellids from Uttar Pradesh I. *Journal of Advanced Zoology*, 20 (2): 106–112.
- Omkar and Pervez, A. (2000). New record of coccinellids from Uttar Pradesh II. *Journal of Advanced Zoology*, 21 (1): 43–47.
- Omkar and Pervez, A. (2004). Predaceous coccinellids in India: predator–prey catalogue. *Oriental Insects*, 38 (1): 27–61.
<https://doi.org/10.1080/00305316.2004.10417373>
- Omkar, Kumar, G. and Sahu, J. (2009). Performance of a predatory ladybird beetle, *Anegleis cardoni* on three aphid species. *European Journal of Entomology*, 106: 565–572.
<https://doi.org/10.14411/eje.2009.071>
- Pal, S. and Bhatt, J. (2018). Feeding efficiency of six beetle species against mustard aphid, *Lipaphis erysimi* (Kalt.) in Uttarakhand, India. *International Journal of Current Microbiology and Applied Sciences*, 7 (4): 1970–1976.
<https://doi.org/10.20546/ijemas.2018.704.226>
- Palial, S., Verma, S. C., Sharma, P. L., Chandel, R. S., Kumar, A., Gupta, M., Sharma, N. and Sharma, P. (2022). Biology, predatory potential and growth parameters of the syrphid fly, *Scaeva pyrastris* (L.) (Diptera: Syrphidae) feeding on the cabbage aphid, *Brevicoryne brassicae* (L.). *Egyptian Journal of Biological Pest Control*, 32: 134.
<https://doi.org/10.1186/s41938-022-00632-5>
- Panchal, K. S., Shetgar, S. S. and Deokar, M. D. (2013). Biology and consumption capacity of *Coccinella septempunctata* linnaeus on aphids of three different crops. *Journal of Entomological Research*, 37 (3): 233–237.
- Pandey, K. P., Kumar, A., Shanker, S. and Tripathi, C. P. M. (1985). First record of *Pachyneuron aphidis* (Bouche) (Pteromalidae: Hymenoptera), a hyperparasitoid of *Diaeretiella rapae* (M'Intosh) (Aphidiidae: Hymenoptera) from India. *Current Science*, 54 (14): 710.
- Pandi, G. G. P., Paul, B., Shah, V. and Shankarganesh, K. (2012). Feeding potential and biology of coccinellid predator *Cheilomenes sexmaculata* (Fabricius) (Coleoptera) on aphid hosts. *Indian Journal of Entomology*, 74 (4): 388–393.

- Parija, S. and Biswal, L. (2023). Seasonal incidence of insect pests and predatory fauna on toria (*Brassica rapa*) in west central Tableland of Odisha, India. *International Journal of Environment and Climate Change*, 13 (11): 231–240.
<https://doi.org/10.9734/ijec/2023/v13i113162>
- Patel, R. A. (2015). Feeding potentiality of *Menochilus sexmaculatus* Fab. on different aphid species. *International Journal on Recent and Innovation Trends in Computing and Communication*, 3 (7): 4426–4430.
- Patel, R. M., Chaudhari, S. J., Rabari, P. H., Patel, B. C. and Dodia, D. A. (2019). Incidence of aphid (*Lipaphis erysimi* Kalt.) in mustard and their fluctuation with biotic and abiotic factors. *Journal of Entomology and Zoology Studies*, 7 (5): 946–950.
- Patel, S. R., Awasthi, A. K. and Tomar, R. K. S. (2004). Assessment of yield losses in mustard (*Brassica juncea* L.) due to mustard aphid (*Lipaphis erysimi* Kalt.) under different thermal environments in eastern central India. *Applied Ecology and Environmental Research*, 2 (1): 1–15.
<https://doi.org/10.15666/aer/02001015>
- Patro, B. (2007). Integrated management of *Lipaphis erysimi* (Kaltenbach) in oilseed *Brassica* crops: a review. *Journal of Oilseed Research*, 24 (1): 1–11.
- Pervez, A. and Chandra, S. (2018). Host plant-mediated prey preference and consumption by an aphidophagous ladybird, *Menochilus sexmaculatus* (Fabricius) (Coleoptera: Coccinellidae). *Egyptian Journal of Biological Pest Control*, 28: 54.
<https://doi.org/10.1186/s41938-018-0060-1>
- Pervez, A. and Kumar, R. (2017). Preference of the aphidophagous ladybird *Propylea dissecta* for two species of aphids reared on toxic host plants. *European Journal of Environmental Sciences*, 7 (2): 130–134.
<https://doi.org/10.14712/23361964.2017.12>
- Pervez, A. and Omkar (2004). Prey-dependent life attributes of an aphidophagous ladybird beetle, *Propylea dissecta*. *Biocontrol Science and Technology*, 14 (4): 385–396.
<https://doi.org/10.1080/09583150410001683547>
- Pervez, A., Omkar, and Harsue, M. A. (2020a). Coccinellids on crops: nature's gift for farmers, In: Chakravarthy, A. K. (Ed.), *Innovative Pest Management Approaches for the 21st century*. Springer Nature Singapore Pte Ltd., pp. 429–460.
- Pervez, A., Nunes, Maia, A. H. N. and Bozdoğan, H. (2020b). Reproduction and demography of an aphidophagous ladybird, *Hippodamia variegata* on six aphid species. *International Journal of Tropical Insect Science*, 40: 541–548.
<https://doi.org/10.1007/s42690-020-00101-2>
- Pervez, A., Yadav, M. and Khan, M. (2020c). Diversity of predaceous coccinellid beetles (Coleoptera: Coccinellidae) in Uttarakhand, North India. *Journal of Mountain Research*, 15: 7–20.
<https://doi.org/10.51220/jmr.v15i1.2>
- Poddar, S. C. and Ghosh, A. K. (1984). Bionomics of an aphidophagous coccinellid predator *Pseudaspidimerus circumflexa* (Motschulsky) (Coccinellidae: Aspidimerini). *Records of the Zoological Survey of India*, 81 (3–4): 67–76.
<https://doi.org/10.26515/rzsi/v81/i34/1984/161219>
- Prabhakar, A. K. and Roy, S. P. (2010). Evaluation of the consumption rates of dominant coccinellid predators on aphids in north– east Bihar. *The Bioscan*, 5 (3): 491–493.
- Pradhan, P. P., Borkakati, R. N. and Saikia, D. K. (2020a). Seasonal incidence of insect pests and natural enemies of mustard in relation to meteorological parameters. *Journal of Entomology and Zoology Studies*, 8 (1): 1538–1542.
- Pradhan, P. P., Borkakati, R. N. and Saikia, D. K. (2020b). Insect pests of mustard and their natural enemies in Assam. *International Journal of Current Microbiology and Applied Sciences*, 9 (7): 2785–2790.
<https://doi.org/10.20546/ijemas.2020.907.328>
- Prakash, S. and Rani, P. (2015). Ecological succession of aphids and their natural enemies of cauliflower. *Research Spectra*, 1 (2–3): 89–90.
- Prodanova, N., Naslednikova, M. and Tarasova, O. (2023). Study of the impact of anthropogenic activities on the environment: problems and prospects of sustainable nature management. *E3S Web of Conferences*, 420: 04001.
<https://doi.org/10.1051/e3sconf/202342004001>
- Rafi, U., Usmani, M. K. and Akhtar, M. S. (2010). Aphids of ornamental plants and winter vegetables and their aphidiine parasitoids (Hymenoptera: Braconidae) in Aligarh region, Uttar Pradesh. *Journal of Threatened Taxa*, 2 (9): 1162–1164.
<https://doi.org/10.11609/JoTT.o2399.1162-4>
- Rahman, K. A. (1940). Important insect predators of India. *Proceedings of the Indian Academy of Sciences*, 12: 67–74.
<https://doi.org/10.1007/BF03049103>
- Raj, D. and Sharma, G. D. (1993). Prevalence of *Diaeretiella rapae* M'Intosh (Hymenoptera: Braconidae) on aphid complex infesting rapeseed. *Journal of Insect Science*, 6: 285–286.
- Rajeshwari, D. and Singh, K. M. (2022). Studies on seasonal incidence of natural enemies of pest complex on broad leaf mustard, *Brassica juncea* var. *rugosa* Roxb. Tsen and Lee in the Valley of Manipur. *Biological Forum – An International Journal*, 14 (3): 1229–1233.
<https://doi.org/10.9734/ijec/2022/v12i930740>
- Rajput, K. (2022). Biological control of mustard aphids: *Diaeretiella rapae* (M'Intosh), a potential biocontrol agent of mustard aphid, *Lipaphis erysimi* Kalt. Lambert Academic Publishing, 72 pp.

- Rakhshani, E., Barahoei, H., Ahmad, Z., Tomanović, Z., Janković, M., Petrović, A., Bodlah, I. and Starý, P. (2012). New species and additional evidence of aphid parasitoids. (Hymenoptera: Braconidae: Aphidiinae) from India. *Zootaxa*, 3397: 45–54.
<https://doi.org/10.11646/zootaxa.3397.1.3>
- Ram, N., Pawariya, V., Mundiya, R. and Bajia, R. (2022). Population dynamics of mustard aphid, *Lipaphis erysimi* (Kalt) in relation to biotic factors. *Journal of Biotechnology and Crop Science*, 10 (15): 183–187.
<https://doi.org/10.5958/25825089.2022.00022.6>
- Ramaseshiah, G., Bhat, K. V. and Dharmadhikari, P. R. (1969). Influence of host aphid, host plant and temperature on the laboratory breeding of *Lysiphlebus testaceipes*. *Indian Journal of Entomology*, 30 (4): 281–285.
- Rana, J. S. (2006). Response of *Coccinella septempunctata* and *Menochilus sexmaculatus* (Coleoptera: Coccinellidae) to their aphid prey, *Lipaphis erysimi* (Hemiptera: Aphididae) in rapeseed–mustard. *European Journal of Entomology*, 103: 81–84.
<https://doi.org/10.14411/eje.2006.013>
- Rao, B. S., Rao, A. S. and Nagalingam, B. (1997). Biology of two coccinellid predators on three species of aphids infesting different crops. *Madras Agricultural Journal*, 84 (2): 53–56.
<https://doi.org/10.29321/MAJ.10.A00836>
- Rao, V. P., Dharmadhikari, P. R., Ramaseshiah, G., Phalak, V. R. and Madhavan, T. V. S. (1969). Study of natural enemies of aphids (for the U.S.A.). *Report of the Commonwealth Institute of Biological Control*, 1968: 45–47
- Rao, V. P., Ramaseshiah, G., Sethumadhavan, T. V. and Ramaswamy, P. (1970). Study of natural enemies of aphids (for the U. S. A.). *Report of the Commonwealth Institute of Biological Control*, 1969: 45–47.
- Ravi, G. B., Reji Rani, O. P., Sudharma, K. and Kavya, M. K. (2014). Insect pests of cabbage and cauliflower and their natural enemies in agro ecosystem of Kerala. *Entomon*, 39 (3): 151–160.
<https://doi.org/10.33307/entomon.v39i3.56>
- Ray, S. K. (1967). Observations on the natural predators of the aphid pests in the Ranchi district with special reference to *Cheilomenes sexmaculata* (Fab.). *Indian Agriculture*, 11: 117–120.
- Raychaudhuri, D. N., Poddar, S. C. and Raychaudhuri, D. (1982). Study of the genus *Aphidius* (Hymenoptera: Aphidiidae) of India. *Entomon*, 7 (1): 11–22.
- Roy, P. and Basu, S. K. (1977). Bionomics of aphidophagous syrphid flies. *Indian Journal of Entomology*, 39 (2): 165–174.
- Sachan, G. C. (1997). Cultural control of aphids: a review and bibliography. *Journal of Aphidology*, 11 (1): 37–40.
<https://doi.org/10.2307/1166744>
- Saha, B. K. (1988). Studies on aphid–aphidiid complex of Jammu and Kashmir. Ph.D. thesis, University of Calcutta, Kolkata, West Bengal, India. 497 pp.
<http://hdl.handle.net/10603/163506>
- Saha, J. L., Poddar, S. C., Das, S. K., Agarwala, B. K. and Raychaudhuri, D. (1982). Studies on the aphid parasites (Hymenoptera: Aphidiidae) from Himachal Pradesh, India. *Akitu NS*, 44: 1–12.
- Saharia, D. (1981). Biology of *Coccinella repanda* Thunb. an important predator of mustard aphid *Lipaphis erysimi* (Kltb.) in Assam. *Journal of Research, Assam Agricultural University*, 2 (1): 66–73.
- Salalia, R. and Isher, A. (2011). On farm validation of biocontrol of mustard aphid (*Lipaphis erysimi*) Kalt. by lady bird beetle (*Coccinella septempunctata*). *Indian Journal of Applied Entomology*, 25 (1): 29–32.
- Samanta, A. K. (1986). Studies on aphidiids (Hymenoptera)-Aphids (Homoptera) of Meghalaya and Arunachal Pradesh. Ph.D. thesis, University of Calcutta, Kolkata, West Bengal, India. 339 pp.
<http://hdl.handle.net/10603/163445>
- Sarkar, S. K. (1991). Studies on aphid–aphidiid complex of Mizoram. Ph.D. thesis, Culcutta University, Kolkata, India. 452 pp.
<http://hdl.handle.net/10603/154725>
- Saxena, R. C. (1981). Natural enemy complex of *Aphis craccivora* Koch. and *Lipaphis erysimi* (Kalt.) in and around Calcutta, West Bengal. *Science and Culture*, 47 (1): 58–60
- Shaikh, M., Pawar, M., Pandit, S. and Mitra, S. (2020). Natural predators of the mustard aphid *Lipaphis erysimi* show differential predation preferences on the aphids feeding on different cole crops. *Indian Journal of Hill Farming*, 33 (2): 322–330.
- Shankarganesh, K., Suroshe, S. S. and Paul, B. (2015). Relative susceptibility of the Bikaner and Delhi populations of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae), and its predator, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), to different insecticides. *Phytoprotection*, 95 (1): 27–31.
<https://doi.org/10.7202/1031955ar>
- Shantibala, S. (1989). Studies on taxonomy and biology of aphidophagous coccinellids (Coleoptera: Coccinellidae) of Manipur and Nagaland. Ph.D. thesis, Manipur University, Canchipur, Imphal, India. 263 pp.
<http://hdl.handle.net/10603/150284>
- Sharma, A. K. and Subba Rao, B. R. (1964). A further contribution to the knowledge of the taxonomy and biology of Aphidiidae (Ichneumonoidea: Hymenoptera) with particular reference to Indian fauna. *Indian Journal of Entomology*, 26 (4): 258–260.

- Sharma, D. K., Varma, G. C. and Kishore, L. (1997). Feeding capacity of predators of mustard aphid, *Lipaphis erysimi*. *Journal of Aphidology*, 11 (2): 171–174.
- Sharma, J. C. (1973). Predation of *Menochilus sexmaculatus* Fab. on different species of aphids. *Indian Journal of Entomology*, 35 (4): 343–344.
- Sharma, P. L., Chauhan, U. and Sharma, K. C. (2015). Studies on the diversity of predatory coccinellid beetles (Coleoptera) in different agro-climatic zones of Himachal Pradesh. *The Bioscan*, 10 (3): 981–985.
- Sharma, P. K. and Joshi, P. C. (2010). New records of coccinellid beetles (Coccinellidae: Coleoptera) from district Dehradun, (Uttarakhand), India. *New York Science Journal*, 3 (6): 112–120.
- Sheileja, T., Maheswari, T. U. and Shantibala, T. (2022). Evaluation of feeding potential and prey preference of *Chrysoperla zastrowi sillemi* (Esben – Peterson) on different Aphid Species. *Biological Forum – An International Journal*, 14 (2a): 188–192.
- Shenhmar, M. and Brar, K. S. (1995). Biological control of mustard aphid, *Lipaphis erysimi* (Kaltenbach) in the Punjab. *Journal of Biological Control*, 9 (1): 9–12.
- Shukla, A. and Kumar, A. (2024). Population dynamics of mustard aphid *Lipaphis erysimi* (Kalt) and its coccinellid predators. *Indian Journal of Entomology*, No. e24553: 1–3. <https://doi.org/10.55446/IJE.2024.1553>
- Shukla, A. N., Tripathi C. P. M. and Singh, R. (1994). Population growth of mustard aphid, *Lipaphis erysimi* Kalt. on different food plants. *Bioved*, 5 (1): 15–20.
- Singh, N. N. and Kumar, M. (2000). Potential of *Chrysoperla carnea* (Stephes) in suppression of mustard aphid population. *Seventh National Symposium on Aphidology*, 11–13 March, 2000, Department of Zoology, D.D.U. Gorakhpur University, Gorakhpur, abstract. p. 35.
- Singh, D. and Singh, H. (1994). Correlation coefficients between abiotic, biotic factors (predators and parasitoid) and mustard aphid, *Lipaphis erysimi* (Kalt.) population on rapeseed – mustard. *Journal of Aphidology*, 8 (1–2): 102–109.
- Singh, G. and Singh, R. (2017). Updated checklist of food plants of Macrosiphini (Aphididae: Hemiptera) in India–2. *International Journal of Research Studies in Zoology*, 3 (1): 42–76. <http://dx.doi.org/10.20431/2454-941X.0301004>
- Singh, K. and Singh N. N. (2013). Preying capacity of different established predators of the aphid *Lipaphis erysimi* (Kalt.) infesting rapeseed–mustard crop in laboratory conditions. *Plant Protection Science*, 49: 84–88. <https://doi.org/10.17221/66/2011-PPS>
- Singh, N. S., Paonam, M. S. and Singh, T. K. (1991). Aphid parasitoids (Hymenoptera: Aphidiidae) from Nagaland, northeast India. *Journal of Aphidology*, 5 (1–2): 61–63.
- Singh, O. P. and Rawat, R. R. (1981). New record of *Diaraetus rapae* M'Intosh as an endoparasite of *Lipaphis erysimi* (Kalt.) from Madhya Pradesh, India. *Science and Culture*, 47: 288–290.
- Singh, P. (2009). Seasonal variation in the species composition of aphids and their ladybeetle predators in mustard crop and rose plantations in relation to chemical changes of the infested plant foliage. Ph.D. thesis, Mahatma Jyotiba Phule Rohilkhand University, Bareilly, Uttar Pradesh, India. 294 pp. <http://hdl.handle.net/10603/325752>
- Singh, P. M. and Singh, T. K. (1986d). Aphid parasitoids (Hymenoptera: Aphidiidae) of Manipur, In: Agarwala, B. K. (Ed.), *Aphidology in India*. pp. 97–99.
- Singh, P. M. (1987). Studies on aphid parasitoids (Hymenoptera: Aphidiidae) of Manipur. Ph.D. thesis, Manipur University, Imphal, Manipur, India. 231 pp. <http://hdl.handle.net/10603/150302>
- Singh, P. M., Devjani, P., Devikarani, K. and Singh, T. K. (2011). Biodiversity, distribution and host range of the genus *Ephedrus* Haliday (Hymenoptera: Aphidiidae) in Manipur, N. E. India. *Journal of Experimental Sciences*, 2 (9): 24–26.
- Singh, R. (2014a). *Elements of Entomology*. Second Edition. Rastogi Publications, Meerut, U. P., India. 492 pp.
- Singh, R. and Singh, G. (2016). Aphids and their biocontrol, In: Omkar (Ed.), *Ecofriendly Pest Management for Food Security*. Academic Press. pp. 63–108. <https://doi.org/10.1016/B978-0-12-803265-7.00003-8>
- Singh, R., Tiwari, A. K. and Tiwari, K. M. (2024a). Checklist of aphidophagous Neuroptera (Insecta: Arthropoda) in India. *International Journal of Biological Innovations*, 6 (2): 99–127. <https://doi.org/10.46505/IJBI.2024.6204>
- Singh, R., Singh, B. B. and Sharma, A. K. (2024b). Checklist of aphidophagous spiders (Araneae: Arachnida: Arthropoda) in India. *Serket*, 20 (3): 288–310.
- Singh, R., Verma, A. K. and Singh, G. (2023). *Food Plant Catalogue of Indian Aphids (Homoptera: Aphididae)*. Asian Biological Research Foundation, Prayagraj, India; Nature Light Publications, Pune, India. 350 pp. <https://doi.org/10.2139/ssrn.4447059>
- Singh, T. K., Devjani, P. and Bijaya, P. (2002). Predator complex of major aphids in cauliflower agro–ecosystem in Manipur. *Journal of Aphidology*, 16 (1–2): 97–102.
- Singh, V. (2014b). Occurrence of insect pests on cabbage and their natural enemies in the periphery of Raipur. *Journal of Applied Zoological Researches*, 25 (2): 145–148.
- Singh, Y. P., Meghwal, H. P. and Singh, S. P. (2009). Evaluation of some bioagents against mustard aphid (*Lipaphis erysimi* (Kaltenbach)) (Hemiptera: Aphididae) on single plant in field condition. *Journal of Biological Control*, 23 (1): 95–97.

- Singh, Y. P. and Meghwal, H. P. (2010). Evaluation of some bioagents against mustard aphid (*Lipaphis erysimi* Kaltenbach) (Homoptera: Aphididae) under field conditions. *Indian Journal of Entomology*, 72 (1): 66–70.
- Singh, Y. P., Meghwal, H. P. and Singh, S. P. (2008). Biology and feeding potential of *Cheilomenes sexmaculata* Fabricius (Coleoptera: Coccinellidae) on mustard aphid. *Annals of Arid Zone*, 47 (2): 185–190.
- Soni, R., Doel, G. S. and Brar, K. S. (2004). Feeding potential of coccinellids on mustard aphid, *Lipaphis erysimi*. (Kalt). *Insect Environment*, 10 (1): 15–16.
- Soni, S., Kumar, S., Sood, A. K. and Rana, R. S. (2021). Modeling of aphid complex and its associated natural enemies in rapeseed–mustard in relation to climatic factors. *Journal of Agrometeorology*, 23 (2): 207–212.
<https://doi.org/10.54386/jam.v23i2.70>
- Srivastava, A. S., Katiyar, R., Upadhyay, K. D. and Singh, S. V. (1982). Studies on the host preference of *Coccinella repanda* Thunberg (Coleoptera: Coccinellidae). *Indian Journal of Entomology*, 49 (4): 551–552.
- Suary, P. and Bhagat, R. C. (1978). Records and descriptions of aphid parasitoids from Kashmir, India (Hymenoptera: Aphidiidae). *Acta Entomologica Bohemoslovaca*, 75 (6): 387–393.
- Suary, P. and Ghosh, A. K. (1975). Aphid parasites (Hymenoptera: Aphidiidae) from Meghalaya, India. *Oriental Insects*, 9 (3): 343–349.
<https://doi.org/10.1080/00305316.1975.10434504>
- Suary, P. and Ghosh, A. K. (1978). Further records of aphid parasitoids (Hymenoptera: Aphidiidae) from Meghalaya, India. *Oriental Insects*, 12 (1): 77–80.
<https://doi.org/10.1080/00305316.1978.10434553>
- Suary, P. and Ghosh, A. K. (1983). Aphid parasitoids of India and adjacent countries (Hymenoptera: Aphidiidae). Zoological Survey of India. Technical Monograph, No. 7: 1–96.
- Subharani, S., Thorat, S. S., Abem, N., Amit Kumar, L. and Singh, T. K. (2010). A database on parasitoid of insect pests of crops in Manipur, India. *The Journal of Plant Protection Sciences*, 2 (2): 66–72.
- Subhrani, S., Singh, M. P. and Singh, T. K. (2006). Biodiversity of aphidiine parasitoids (Hymenoptera: Braconidae) on certain crops in Manipur. *Journal of Aphidology*, 20 (1–2): 1–4.
- Tamili, H. K. (1988). Taxonomy of Aphidiids – Aphids of Sikkim and Hilly areas of West Bengal. Ph.D. thesis, University of Calcutta, Kolkata, West Bengal, India. 370 pp.
<http://hdl.handle.net/10603/163376>
- Thamilvel, D. (2009). Population dynamics and management of aphids in vegetable ecosystem. Ph.D. thesis, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala. 266 pp.
<http://14.139.185.57:8080/jspui/bitstream/123456789/8932/1/172968.pdf>
- Tiwari, K. M., Singh, R. and Tripathi, P. N. (2024a). Biosystematics, economic importance, natural enemies and food plants of *Lipaphis* (*Lipaphis*) *erysimi* (Kaltenbach, 1843) (Hemiptera: Aphididae) in India. *Munis Entomology and Zoology*, 19 (2): 615–635.
- Tiwari, K. M., Tripathi, P. N. and Singh R. (2024b). Biodiversity of aphidophagous predators in Ayodhya district, Uttar Pradesh. *Arthropods*, 13 (3): 120–156.
- Varmora, J. M., Raghvani, K. L., Joshi, M. D., Makadia, R. R., Dalwadi, N. G. and Boricha, H. V. (2009). Population dynamics of aphid, *Lipaphis erysimi* (Kalt.) and predators on cabbage. *Asian Sciences*, 4: 56–58.
- Veda, O. P. and Shaw, S. S. (1988). Population dynamics of mustard aphids, *Lipaphis erysimi* (Kalt.) and its parasite, *Diaretus rapae* (M'Intosh). *Agricultural Science Digest, Karnal*, 8: 189–190.
- Verma, P. K., Varma, A. and Aggarwal, G. (2023). Studies on insect–pests complex of rapeseed mustard and their natural species in Roorkee Region, Uttarakhand, India. *Biological Forum – An International Journal*, 15 (5): 529–533.
- Verma, S. N., Gargav, V. P. and Mittal, S. (1983). Host preference of six spotted ladybird beetle, *Menochilus sexmaculatus* (Fabr.). *Indian Journal of Plant Protection*, 11: 66–69.
- Waluniba, W., Ao, M. A., Singh, H. K., Neog, P. and Lytan, D. (2016). Bio–diversity of insect pests and its natural enemies in cabbage ecosystem of Nagaland, northeast India. *Environment and Ecology*, 34 (3B): 1346–1350.
- WFO (2024). The World Flora Online. <https://www.worldfloraonline.org> (Accessed 5 August 2024).
- WSC. (2024). World Spider Catalog. Version 25. Natural History Museum Bern, online at <http://wsc.nmbe.ch> (Accessed 5 August 2024).
<https://doi.org/10.24436/2>
- Yadav, R. V., Kumari, V. and Saxena, R. (2023). Population dynamics of aphid and their natural enemies in mustard based on meteorological parameters using principal component analysis. *Journal of Agrometeorology*, 25 (3): 458–461.
<https://doi.org/10.54386/jam.v25i3.2209>
- Zalavadia, A. G. and Kapadia, M. N. (2000). Comparative predation capacity of *Coccinella septempunctata* L. on three prey aphids. *Seventh National Symposium on Aphidology*, 11–13 March, 2000, Department of Zoology, D.D.U. Gorakhpur University, Gorakhpur, abstract. p. 49.